

Finite field Nullstellensatz and Grassmannians

E. Ballico

Dipartimento di Matematica
Università di Trento
38050 Povo (TN), Italy
ballico@science.unitn.it

A. Cossidente

Dipartimento di Matematica
Università della Basilicata
via N. Sauro, 85
85100 Potenza, Italy
cossidente@unibas.it

Abstract

Let $X \subset \mathbb{P}^N$ be a projective variety defined over the Galois field $GF(q)$. Denote by $X(q)$ the set of $GF(q)$ -rational points of X . Let k be an integer. We say that the pair $(X, X(q))$ satisfies the *Finite Field Nullstellensatz of order k* , (the $FFN(k)$, for short), if every homogeneous form of degree $\leq k$ on $\mathbb{P}^N(K)$ vanishing on $X(q)$, vanishes on $X(K)$. Here, we prove the Finite Field Nullstellensatz $FFN(q)$ for any Grassmann variety.

1 Introduction

Let p be a prime and q be a power of p . Let $GF(q)$ be the Galois field of order q and let K denote the algebraic closure of $GF(q)$. Let $X \subset \mathbb{P}^N$ be a projective variety defined over $GF(q)$. Let $X(q)$ (resp. $X(K)$) denote the set of all $GF(q)$ -rational points (resp. the K -rational points) of X . Then $X(q)$ is a finite subset of \mathbb{P}^N . We will denote the N -dimensional projective space over $GF(q)$ by $\mathbb{P}^N(q)$.

Let k be an integer. We say that the pair $(X, X(q))$ satisfies the *Finite Field Nullstellensatz of order k* , (the $FFN(k)$, for short), if every homogeneous form of degree $\leq k$ on $\mathbb{P}^N(K)$ vanishing on $X(q)$, vanishes on $X(K)$.

When X is a geometrically interesting variety, the homogeneous ideal of X (i.e. the homogeneous ideal of $X(K)$) is often known, and the knowledge of $FFN(k)$ gives a complete description, up to degree k , of the homogeneous ideal of $X(q)$.

In this note, we will consider the case in which X is a *Grassmann variety*, and the embedding $X \subset \mathbb{P}^N$ is given by the Plücker embedding.

Recall that any Grassmannian, say G , is defined over $GF(p)$, and that the Plücker embedding of G is defined over $GF(p)$. Hence, both are defined over $GF(q)$, and so we may consider $FFN(k)$ for the pair $(G, G(q))$ with respect to the Plücker embedding. For more details on Grassmannians, see [5], [6], [7].

The aim of this note is to prove the following theorem.

Theorem 1.1 *Let $G \subset \mathbb{P}^N$ be the Plücker embedding of any Grassmannian. Then the pair $(G, G(q))$ satisfies $FFN(q)$.*

Remark 1.2 Theorem 1.1 is sharp because $\mathbb{P}^N(q)$ is the union of the $GF(q)$ -rational points of $q + 1$ hyperplanes defined over $GF(q)$. Hence, for any subvariety X of \mathbb{P}^N defined over $GF(q)$, the pair $(X, X(q))$ never has property $FFN(q + 1)$.

In dealing with Grassmann spaces, Moorhouse in [4, Theorem 2] proved that $FFN(q - 1)$ is true for the Plücker embedding of any Grassmannian. Our theorem is stronger than Moorhouse's result and the approach is different. In particular, [4, Theorem 2] is a consequence of Theorem 1.1. In the case of the Klein quadric, that is, for the Plücker embedding of the Grassmannian of lines of \mathbb{P}^3 , see [2]. For other results on the Finite Field Nullstellensatz, see [1].

Over the last decade, there have been numerous papers, too many to be quoted here, dealing with the p -rank of incidence matrices of classes of incidence structures, such as projective spaces, translation planes, orthogonal spaces, generalized quadrangles, unitals, designs, Hermitian varieties, Grassmann varieties. Our result is a contribution to current research on p -ranks; nevertheless, we consider our techniques a significant step toward the FFN problem for other varieties, which we feel has an independent interest.

2 The Proof of Theorem 1.1

For any integers x, n , with $0 \leq x < n$, let $G(x, n)$ be the Grassmannian of x -dimensional projective linear subspaces of \mathbb{P}^n . Hence $\dim(G(x, n)) = (n - x)(x + 1)$. It turns out that $G(0, n) = \mathbb{P}^n$ and $G(n - 1, n) = \mathbb{P}^{n*} \cong \mathbb{P}^n$.

Set $N = N(x, n) := ((n + 1)! / ((n - x)!(x + 1)!)) - 1$. The Plücker embedding of $G(x, n)$ embeds $G(x, n)$ into \mathbb{P}^N .

Let X_0, \dots, X_n be homogeneous coordinates in \mathbb{P}^n .

We will prove Theorem 1 for $G(x, n)$ by induction on n . The case $x = 0$ just means that any homogeneous polynomial $f(X_0, \dots, X_n)$ with $\deg(f) \leq q$ and with f vanishing at each point of $\mathbb{P}^n(q)$, is identically zero. Hence the case $x = 0$ is trivially true for any n . Now, we fix the integer x with $0 < x \leq n - 2$, and we assume the result holds true for the pair $(n - 1, x)$. We regard $G(x, n)$ as embedded in \mathbb{P}^N by the Plücker embedding. Fix an integer $t \leq q$ and let $f \in K[X_0, \dots, X_n]$ be a homogeneous polynomial with $\deg(f) = t$ and $f(P) = 0$ for every point $P \in G(x, n)(q)$. The projective space $\mathbb{P}^n(q)$ has $a := (q^{n+1} - 1) / (q - 1)$ hyperplanes, and each of them defines an embedding of $G(x, n - 1)(q)$ into $G(x, n)(q)$, and of $G(x, n - 1)(K)$ into $G(x, n)(K)$. Let $\{H_i\}$, $1 \leq i \leq a$, be the set of all hyperplanes

of $\mathbb{P}^n(q)$ and $\alpha_i : G(x, n-1) \rightarrow G(x, n)$ the associated embedding. Notice that the restriction to $G(x, n-1)$ of the Plücker embedding of $G(x, n)$ induces the Plücker embedding of $G(x, n-1)$. Hence, by the inductive assumption, we may assume that for every $1 \leq i \leq a$, f vanishes on every point of $\alpha_i(G(x, n-1))(K)$. It is sufficient to prove that f vanishes at a general $Q \in G(x, n)(K)$.

Any such general point Q is contained in a line $D \subset G(x, n) \subset \mathbb{P}^N$ defined over K and corresponds to fixing a $(x+1)$ -dimensional linear subspace V of \mathbb{P}^n and taking all x -dimensional linear subspaces of V . We choose a general such line D . By the generality of Q and of D , we may assume that $V(K)$ is a general $(x+1)$ -dimensional linear subspace of $\mathbb{P}^n(K)$. Hence, we may assume that $V(K)$ intersects transversally each hyperplane $H_i(K)$, $1 \leq i \leq a$, that is, $\dim(V(K) \cap H_i(K)) = x$, for every $1 \leq i \leq a$, and that $V(K) \cap H_i(K) \neq V(K) \cap H_j(K)$, for $i \neq j$. Thus D intersects each $\alpha_i(G(x-1, n)(K))$ at a different point. Since $\deg(f) = t \leq q < a$, we obtain $f|_D \cong 0$. Since $Q \in D$, we obtain $f(Q) = 0$, as wanted. Now, we consider the case $x = n-1$. We have $G(n-1, n) = \mathbb{P}^{n*} \cong \mathbb{P}^n$, and the proof is as for the case $x = 0$.

Acknowledgement. This research was carried out within the activity of G.N.S.A.G.A. of the Italian C.N.R. with the support of the Italian Ministry for Research and Technology.

References

- [1] E. Ballico and A. Cossidente, On the finite field Nullstellensatz, *Australas. J. Combin.* **21** (2000), 57–60.
- [2] A. Blokhuis and G.E. Moorhouse, Some p -ranks related to orthogonal spaces, *J. Algebraic Combin.* **4** (1995), 295–316.
- [3] G.E. Moorhouse, Some p -ranks related to Hermitian varieties, *J. Stat. Plan. Inf.* **56** (1996), 229–241
- [4] G.E. Moorhouse, Some p -ranks related to finite geometric structures, in *Mostly Finite Geometries*, ed. N.L. Johnson, Marcel Dekker Inc., 1997, 353–364.
- [5] P. Griffiths and J. Harris, *Principles of Algebraic Geometry*, Wiley and Sons, 1978.
- [6] J. Harris, *Algebraic Geometry, A First Course*, Springer-Verlag, New York, 1992.
- [7] J.W.P. Hirschfeld and J.A. Thas, *General Galois Geometries*, Oxford University Press, Oxford, 1991.

(Received 17/11/2000)

