

Further results on proper and strong set colorings of graphs

S.M. HEGDE

*Department of Mathematical and Computational Sciences
National Institute of Technology Karnataka, Surathkal
Srinivasnagar - 575 025 Mangalore
India
smhegde@nitk.ac.in*

M.K. SUMANA

*Department of Mathematics
Dr. N.S.A.M. First Grade College, Nitte
Karkala - 574 110 Udupi
India
sgbarlaya@gmail.com*

Abstract

A *set coloring* α of a graph G is defined as an assignment of distinct subsets of a finite set X of *colors* to the vertices of G such that all the colors of the edges which are obtained as the symmetric differences of the sets assigned to their end-vertices are distinct. Additionally, if all the sets on the vertices and edges of G form the set of all nonempty subsets of X , then the coloring α is said to be a *strong set coloring*, and the graph G is called *strongly set colorable*. If all the nonempty subsets of X are obtained on the edges of G , then α is called a *proper set coloring*, and such a graph G is called *properly set colorable*. The *set coloring number* of a graph G , denoted by $\sigma(G)$, is the smallest cardinality of a set X such that G has a set coloring with respect to X .

This paper discusses the set coloring number of certain classes of graphs and the construction of strongly set colorable caterpillars which are also properly set colorable. An upper bound for b is found for $K_{3,b}$ to admit set coloring with set coloring number n .

1 Introduction

For all standard notation and terminology in graph theory we follow Harary [4] and West [7]. In this paper we consider only finite simple graphs.

The notion of set coloring of a graph was introduced by Hegde [5] in 2009. Acharya [1] initiated a general study of labeling of the vertices and the edges of a graph using subsets of a set and indicated their potential application in a variety of other areas of human enquiry. Given a graph $G = (V, E)$ and a nonempty set X of n colors, a function $f : V \rightarrow 2^X$ can be defined as the assignment of the colors $f(v)$, to each of the vertices $v \in V$, and given such a function f on the vertex set V , we define $f^\oplus : E \rightarrow 2^X$ which assigns colors to the edges $e = uv \in E$ as $f^\oplus(e) = f(u) \oplus f(v)$.

A graph G is said to be a *set colorable* graph if both f and f^\oplus are injective. A graph G is said to be *properly set colorable* if it is set colorable with $f^\oplus(E) = 2^X \setminus \emptyset$, and G is said to be *strongly set colorable* if $f(V) \cup f^\oplus(E) = 2^X \setminus \emptyset$ and $f(V) \cap f^\oplus(E) = \emptyset$.

2 Set coloring number

The *set coloring number* $\sigma(G)$ of a graph G is the least cardinality of a set X with respect to which G has a set coloring.

For any graph $G(V, E)$, we have $\lceil \log_2\{|E| + 1\} \rceil \leq \sigma(G) \leq |V| - 1$, and the bounds are best possible as mentioned in Hegde [5]. In this section we find the set coloring number of some classes of cycles and complete bipartite graphs.

Theorem 1 *Given any positive integer $n \geq 3$, $\sigma(C_{2^n-3}) = n + 1 = \sigma(C_{2^n-2})$.*

Proof: Let the vertices of C_{2^n-3} be denoted by $v_1, v_2, \dots, v_{2^n-3}$ such that $(v_i, v_{i+1}) \in E$ for all $1 \leq i \leq 2^n - 3$; then $\sigma(C_{2^n-3}) \geq n$. Let us assume that there exist a set coloring (f, f^\oplus) of C_{2^n-3} with respect to a set X of $|X| = n$. Then since both f and f^\oplus are injective, the edges have distinct colors. That is,

$$f^\oplus(v_1, v_2) \oplus f^\oplus(v_2, v_3) \oplus \dots \oplus f^\oplus(v_{2^n-3}, v_1) = \emptyset.$$

Let $A_1, A_2, \dots, A_{2^n-1}$ be the distinct nonempty subsets of X . Assign $A_i = f^\oplus(v_i, v_{i+1})$ for $i = 1, 2, \dots, 2^n - 3$; then we get

$$A_1 \oplus A_2 \oplus \dots \oplus A_{2^n-3} = \emptyset. \quad (1)$$

But we know

$$A_1 \oplus A_2 \oplus \dots \oplus A_{2^n-3} \oplus A_{2^n-2} \oplus A_{2^n-1} = \emptyset. \quad (2)$$

From (1) and (2) we get $A_{2^n-2} \oplus A_{2^n-1} = \emptyset \Rightarrow A_{2^n-2} = A_{2^n-1}$, which is a contradiction. Therefore $\sigma(C_{2^n-3}) > n$.

When $|X| = n + 1$, by using the algorithm given in Molard and Payan [6], the subsets of X are assigned to the vertices of C_{2^n-3} to get the set coloring of C_{2^n-3} .

One can prove by using the same argument that $\sigma(C_{2^n-2}) = n + 1$. \square

Theorem 2 *Given any positive integer $n \geq 3$, $\sigma(K_{3,2^{n-2}+1}) = n + 1$.*

Proof: Let $\sigma(K_{3,b}) = n$. We shall prove that the maximum value that b can have is 2^{n-2} . Let V_1 (containing three vertices) and V_2 be the partition of the vertex set V , and let A_1, A_2, \dots, A_{2^n} be the subsets of a nonempty set X of cardinality n . Assign the sets A_1, A_2, A_3 to the vertices of V_1 under the mapping (f, f^\oplus) defined for the sets $V(G)$ and $E(G)$ respectively.

If $A_1 \oplus A_2 = \{x_1, x_2, \dots, x_k\}$, then let $x_1, x_2, \dots, x_j \in A_1$ and $x_{j+1}, x_{j+2}, \dots, x_k \in A_2$. If A_k is any other set containing $\{x_1, x_2, \dots, x_{k-1}\}$ and A_r is a set containing x_k such that $A_k - \{x_1, x_2, \dots, x_{k-1}\} = A_r - \{x_k\}$, then $A_1 \oplus A_k = A_2 \oplus A_r$, which is a contradiction. Therefore, those sets containing x_k cannot be considered to assign the vertices of V_2 and the number of subsets of X not containing x_k are 2^{n-1} out of which A_1 is assigned to a vertex of V_1 . Therefore the possible number of subsets that can be considered to assign the vertices of V_2 is $2^{n-1} - 1$. Again, if $A_2 \oplus A_3 = \{x'_1, x'_2, \dots, x'_m\}$, then by the similar argument mentioned above, sets containing one of x'_j , $j = 1, 2, \dots, m$, cannot be considered for the assignment to the vertices of V_2 . The number of subsets out of $2^{n-1} - 1$ not containing x'_j is 2^{n-2} , out of which one set (A_2 or A_3) is assigned to a vertex of V_1 . Since the set $A_1 \oplus A_2 \oplus A_3$ can be used for the assignment to a vertex of V_2 , the maximum number of sets that are used for the assignment to the vertices of V_2 is 2^{n-2} . Thus $\sigma(K_{3,2^{n-2}}) = n$. Hence $\sigma(K_{3,2^{n-2}+1}) = n + 1$. \square

3 Properly and strongly set colorable graphs

In Hegde [5], it has been mentioned that a necessary condition for a (p, q) -graph G to be properly set colorable is that $q + 1 = 2^m$ for the positive integer $m = |X|$. This condition says that cycles of length not equal to $2^m - 1$ are not properly set colorable.

Molard and Payan [6] proved the following theorem, which says that the cycles of length $2^m - 1$ are properly set colorable.

Theorem 3 *For every integer $n \geq 2$, it is possible to label the vertices of the cycle C_{2^n-1} by all non zero vectors of the vector space $GF^n(2)$ of dimension n over the finite Galois field $GF(2)$ such that the vectors $x \oplus y$ with x and y adjacent, are also all distinct non-zero vectors of $GF^n(2)$, where \oplus denotes addition in $GF^n(2)$.*

Using this result we shall prove the following.

Theorem 4 *For any two positive integers k_1 and k_2 , $C_{2^{k_1}-1} + C_{2^{k_2}-1}$ is properly set colorable.*

Proof: Consider $G = C_{2^{k_1}-1} + C_{2^{k_2}-1}$ where k_1, k_2 are integers. Then the total number of edges in G is $2^{k_1} - 1 + 2^{k_2} - 1 + (2^{k_1} - 1)(2^{k_2} - 1) = 2^{k_1+k_2} - 1$. By

taking $k_1 + k_2 = m$, the necessary conditions for the existence of proper set coloring is satisfied.

Let $GF^m(2)$ be a vector space over a finite Galois field $GF(2)$, and let $P(x)$ and $Q(y)$ be two primitive irreducible polynomials of degree k_1 and k_2 respectively. Then $\{x^i \text{ mod } P(x)\}$ and $\{y^j \text{ mod } Q(y)\}$ are two disjoint sets of non-zero polynomials of degree at the most $2^{k_1} - 1$ and $2^{k_2} - 1$, respectively, over $GF(2)$. Then label the vertices of the cycles $C_{2^{k_1}-1}$ and $C_{2^{k_2}-1}$ clockwise respectively by $x^1, x^2, \dots, x^{2^{k_1}-1}$ and $y^1, y^2, \dots, y^{2^{k_2}-1}$, where $x^i \neq y^j$ as explained in Mollard and Payan [6].

Analogously we assign the subsets of a nonempty set X of cardinality m to the vertices of G in place of polynomials as follows:

Let $X_1 = \{x_1, x_2, \dots, x_{k_1-1}\}$ and $X_2 = \{y_1, y_2, \dots, y_{k_2-1}\}$ be two disjoint nonempty subsets of X where $X_1 \cup X_2 = X$. The term x^i in each polynomial $\{x^i \text{ mod } P(x)\}$ is replaced by an element x_i of X_1 for $i = 0, 1, 2, \dots, k_1 - 1$ and y^j in each polynomial $\{y^j \text{ mod } Q(y)\}$ is replaced by y_j of X_2 for $j = 1, 2, \dots, k_2 - 1$. Then the polynomials $\{x^i \text{ mod } P(x)\}$ are replaced by the subsets of X_1 by replacing each term of the polynomials by the respective element in X_1 . For instance if $x = x_1$ and $x^2 = x_2$ then $x + x^2 = \{x_1, x_2\}$, and so on. Similarly, the polynomials $\{y^j \text{ mod } Q(y)\}$ are replaced by the subsets of X_2 . One can easily observe that G is properly set colorable. \square

For example, a proper set coloring of $C_{2^3-1} + C_{2^2-1}$ is displayed in Figure 1.




Figure 1: Properly set colored $C_7 + C_3$.

Definition 1 (Gallian [3]) Umbrella graphs are the graphs obtained by joining an edge to the center of a wheel.

From the necessary condition mentioned in Hegde [5], it is clear that no wheel is properly set colorable. The following result shows that the umbrella graphs of the form U_{2^n-1} are properly set colorable.

Corollary 4.1 *An umbrella graph of the form U_{2^n-1} is properly set colorable.*

Proof: Consider the wheel of the form W_{2^n-1} , where n is the cardinality of a set X . Let the pendant vertex w be joined to the central vertex v of the wheel.

For any $x \notin X$, let $X' = X \cup \{x\}$. Assign the set X' to the central vertex, and \emptyset to the pendant vertex w . According to the method explained in Theorem 4, the vertices of the outer cycle C_{2^n-1} are assigned by the nonempty subsets of X . One can easily verify that U_{2^n-1} is properly set colorable. \square

Figure 2 is an example of a proper set coloring of U_7 .




Figure 2: Properly set colored U_7 .

As mentioned in Hegde [5], a necessary condition for a (p, q) graph G to be strongly set colorable is that $p + q + 1 = 2^m$ for the positive integer $m = |X|$. Given below is a relation between strongly set colorable trees and properly set colorable trees.

We call a set having odd number of elements an *odd set* and having even number of elements an *even set*.

Theorem 5 a) *Every properly set colorable tree is strongly set colorable.*
 b) *A strongly set colorable tree whose vertex labels are odd sets is properly set colorable.*

Proof: a) Let T be a properly set colorable tree with a proper set coloring f with respect to a nonempty set X of cardinality m . Let $X' = X \cup \{x\}$. Since

T is properly set colorable, all the subsets of X are assigned to the vertices and $f(T) = \{f(v) \mid v \in V(T)\} = 2^X$ and $f^\oplus(T) = \{f(e) \mid e \in E(T)\} = 2^X - \emptyset$.

Define a function $F : V(T) \rightarrow 2^{X'}$ by $F(v) = f(v) \cup \{x\}$, for all $v \in V(T)$.

Since f and f^\oplus are injective, F and F^\oplus are also injective. Also $F(V(T)) \cap F^\oplus(E(T)) = \emptyset$.

Since $f(T) = 2^X$ and $f^\oplus(T) = 2^X - \emptyset$, we get $F(T) = 2^{X'} - 2^X$ and $F^\oplus(T) = 2^X - \emptyset$. That is, $f^\oplus(T) = F^\oplus(T)$. Further,

$$\begin{aligned} |F(T)| &= 2^{|X'|} - 2^{|X|} \\ &= 2^{m+1} - 2^m \\ &= 2^m(2 - 1) \\ &= 2^m \quad \text{and} \quad |F^\oplus(T)| = 2^m - 1. \end{aligned}$$

Therefore $|F(T)| + |F^\oplus(T)| = 2^m + 2^m - 1 = 2^{m+1} - 1 = 2^{|X'|} - 1$. This implies that F is a strong set coloring of T .

b) Let f be a strong set coloring of T with respect to the set X of n colors, such that $f(v)$ is an odd set for each $v \in V(T)$.

Define $F : V(T) \cup E(T) \rightarrow 2^{X'}$ where $X' = X - \{w\}$, $w \in X$ as

$$F(v) = \begin{cases} f(v) - \{w\} & \text{if } w \in f(v) \\ f(v) & \text{if } w \notin f(v) \\ \emptyset & \text{if } w = f(v). \end{cases}$$

Then we have $F^\oplus(uv) = f^\oplus(uv) - \{w\}$ when $w \in f(u)$ or $w \in f(v)$, and $F^\oplus(uv) = f^\oplus(uv)$ when $w \in f(u)$ and $w \in f(v)$, or $w \notin f(u)$ and $w \notin f(v)$.

This implies that F^\oplus is injective. Further, if $F(v) = f(v) - \{w\}$ then $F(v)$ is an even set and if $F(u) = f(u)$ then $F(u)$ is an odd set, so that $F^\oplus(uv)$ becomes an odd set. The number of even subsets in $f(T)$ is $2^{n-1} - 1$, out of which 2^{n-2} subsets contain w . Therefore the number of odd sets on the edges under F is 2^{n-2} . The remaining $2^{n-2} - 1$ subsets are even sets on the edges under F . Hence the total number of subsets assigned to the edges of T is $2^{n-2} + 2^{n-2} - 1 = 2^{n-1} - 1 = 2^{|X'|} - 1$. This implies that all the nonempty subsets of X' are on the edges of T . Hence F is a proper set coloring of T . \square

Figure 3 illustrates that a properly set colorable tree is strongly set colorable and Figure 4 illustrates that a strongly set colorable tree with odd sets assigned to the vertices is properly set colorable.

Definition 2 (Chen [2]) An (n, k) -banana tree, denoted by $B(n, k)$ is a graph obtained by connecting one leaf of each of n copies of a k -star graph with a single root vertex that is distinct from all the stars.

Theorem 6 For any positive integer, $r \geq 2$, a banana tree $B(k, n)$, where $k = 2^{r/2} - 1$ and $n = 2^{r/2}$, is properly set colorable.




Figure 3: An illustrative example of Theorem 5(a).




Figure 4: An illustrative example of Theorem 5(b).

Proof: Let w_0 be the central vertex, and w_1, w_2, \dots, w_k be the roots of the stars joining the central vertex. Let $v_{i,1}, v_{i,2}, \dots, v_{i,n}$ denote the pendant vertices joining w_i where $i = 1, 2, \dots, k$. Let X be a nonempty set with $|X| = r$ and let X' be a subset of X where $|X'| = r/2$.

We define a mapping $f : V(B) \cup E(B) \rightarrow 2^X$ as follows:

$$f(w_0) = \emptyset, \quad f(w_j) = A_j, \text{ where } A_j \subseteq X', \quad f(v_{i,1}) = B_i, \text{ where } B_i \subseteq X - X', i = 1, 2, \dots, k, \quad f(v_{i,j}) = B_i \cup A_j.$$


Since the A_j and B_i are disjoint, vertices are assigned by the distinct subsets. Therefore the mapping f is injective. Further,

$$f^\oplus(w_i, v_{i,1}) = A_j \oplus B_i, \quad f^\oplus(w_0, w_j) = A_j, \quad f^\oplus(w_j, v_{i,j}) = A_j \oplus (B_i \cup A_j) = B_i$$

which shows f^\oplus is injective. One can see that the total number of edges is $kn + k = (2^{r/2} - 1)(2^{r/2} + 1) = 2^r - 1$, which is equal to the total number of nonempty subsets of X .

Hence $B(k, n)$ is properly set colorable, and by Theorem 5, $B(k, n)$ is also strongly set colorable. \square

Figure 5 is a properly set colored banana tree $B(3, 4)$.

Figure 5: Properly set colored banana tree $B(3, 4)$.

Construction of strongly (properly) set colorable caterpillars:

Construction of an infinite family of strongly (properly) set colorable caterpillars is given below.

Let X_1 be a nonempty set with $|X_1| = m_1$ where $m_1 \geq 2$ is a positive integer. Consider the star $K_{1,2^{m_1-1}-1} = T_0(m_1)$, say. Let u_0 be the central vertex and $v_{11}, v_{12}, \dots, v_{1,2^{m_1-1}-1}$ be the pendant vertices of $T_0(m_1)$. We define a mapping $f_1 : V(T_0(m_1)) \rightarrow 2^{X_1}$ as follows:

$$f_1(u_0) = \{x_0\}, \text{ where } x_0 \in X_1,$$

$$f_1(v_{1,i}) = A_i, \text{ where } A_i \subset X_1 - \{x_0\}, i = 1, 2, \dots, 2^{m_1-1} - 2,$$

$$f_1(v_{1,2^{m_1-1}-1}) = X_1.$$

Clearly f_1, f_1^\oplus are injectives. Let X_2 be a set of cardinality m_2 where $m_2 > m_1$. Introduce new vertices, say $u_{11}, u_{12}, u_{13}, \dots, u_{1k_1}$, where $k_1 = 2^{m_2-1} - 2^{m_1-1}$, and join each of them to $v_{1,2^{m_1-1}-1}$. Let the resulting caterpillar be denoted by $T_1(m_2)$ and define the mapping $f_2 : V(T_1(m_2)) \rightarrow 2^{X_2}$ as follows:

$$\left\{ \begin{array}{l} f_2(u_{1,j}) = B_j \text{ where } B_j \subset X_2 - X_1, \quad j = 1, 2, \dots, 2^{m_2-m_1}; \\ f_2(u_{1,j}) = A_i \cup B_j \text{ where } B_j \subset X_2 - X_1 \\ \quad \text{and } A_i \subset X_1 \setminus \emptyset \text{ containing } \leq (m_1 - 1)/2 \text{ elements if } m_1 \text{ is odd,} \\ \quad \leq (m_1 - 2)/2 \text{ elements if } m_1 \text{ is even and} \\ \quad \text{containing } m_1/2 \text{ elements, where } A_i^c \text{ are not included;} \\ f_2(u_{1,k_1}) = X_2. \end{array} \right.$$

Let $f_2^\oplus : E(T_1(m_2)) \rightarrow 2^{X_2}$ denote the induced edge function defined by $f_2^\oplus(uv) = f_2(u) \oplus f_2(v)$. Then it is not hard to verify that $f_2 \cup f_2^\oplus$ is the extension of strong set coloring of $f_1 \cup f_1^\oplus$ in $T_0(m_1)$ and is also strongly set colorable. Next, introduce $2^{m_3-1} - 2^{m_2-1}$ new vertices, say $v_{21}, v_{22}, \dots, v_{2k_2}$, where $k_2 = 2^{m_3-1} - 2^{m_2-1}$, and join

each of them to u_{1k_1} . Let X_3 be the set of cardinality m_3 where $X_1 \subset X_2 \subset X_3$ and $m_1 < m_2 < m_3$. Let the resulting caterpillar be $T_2(m_3)$.

Define $f_3 : T_2(m_3) \rightarrow 2^{X_3}$ by

$$\left\{ \begin{array}{l} f_3(v_{2,i}) = B_i \text{ where } B_i \subset X_3 - X_2; \\ \quad = A_k \cup B_i \text{ where } B_i \subset X_3 - X_2 \\ \quad \text{and } A_k \subset X_2 \setminus \emptyset \text{ containing } \leq (m_2 - 1)/2 \text{ elements if } m_2 \text{ is odd,} \\ \quad \leq (m_2 - 2)/2 \text{ elements if } m_2 \text{ is even, and} \\ \quad \text{containing } m_2/2 \text{ elements, where } A_k^c \text{ are not included;} \\ f_3(v_{2,k_2}) = X_3. \end{array} \right.$$

It turns out that $f_3 \cup f_3^\oplus$ is the extension of the strongly set colorable mapping $f_2 \cup f_2^\oplus$ of $T_1(m_1)$ to $T_2(m_3)$ which is also strongly set colorable. We may iterate this procedure indefinitely to obtain the strongly set colorable caterpillar at the n^{th} step, $n = 1, 2, \dots$ where $X_1 \subset X_2 \subset \dots \subset X_n$, and $m_1 < m_2 < \dots < m_n$ are chosen quite arbitrarily.

An illustrative example for the construction of strongly set colorable caterpillars is given in Figure 6.




Figure 6: Strongly set colored caterpillar.

Given below is a method of constructing a properly set colorable caterpillar from the strongly set colorable caterpillar constructed above.

Remove an element from X_1 , say x_0 . Let the resulting set be X_1' . The mapping F_1 is defined as follows:

$F_1(u_0) = \emptyset$; $F_1(v_{1,i}) = A_r$ where $A_r \subset X'_1$, $i = 1, 2, \dots, 2^{m_1-1}-2$; $F_1(v_{1,2^{m_1-1}-1}) = X'_1$. Clearly F_1 is injective and since \emptyset is assigned to the central vertex of $T_0(m_1)$, those subsets, namely A_r , assigned to the other end vertices of the edges incident to the central vertex that is $v_{1,i}$, are also on the edges. Therefore $F^\oplus : E(T_0(m_1)) \rightarrow 2^{X'_1}$ is injective. The number of vertices other than the central vertex is $2^{m_1-1} - 1$ and is equal to the number of nonempty subsets of X'_1 . This implies that F_1 is the proper set coloring of $T_0(m_1)$.

Let X'_2 be the set obtained by deleting the same element x_0 from X_2 . The mapping F_2 is defined as follows:

$$\begin{cases} F_2(u_{1,j}) &= B_j \text{ where } B_j \subseteq X'_2 - X'_1; \\ &= A_i \cup B_j, \text{ where } A_i \text{ is the subset of } X'_1 \\ &\quad \text{and } B_j \text{ is the subset of } X'_2 - X'_1; \\ F_2(u_{1,k_1}) &= X'_2. \end{cases}$$

It is not difficult to verify that the mapping F_2 and the induced edge mapping $F_2^\oplus : E(T_1(m_2)) \rightarrow 2^{X'_2}$ are injective. Further, as explained above, $E(T_2(m_1)) = Y(X'_2)$. This implies that F_2 is a proper set coloring of $T_1(m_2)$.

By proceeding in the same manner, we can show that the above constructed strongly set colored caterpillar is properly set colored.




Figure 7: Properly set colored caterpillar.

Acknowledgements

We are thankful to the referees for their useful suggestions/comments in the improvement of the paper.

References

- [1] B.D. Acharya, Set valuations of a graph and their applications, *MRI Lecturer Notes in Applied Mathematics* No. 2, MRI, Allahabad, 1983.
- [2] W. C. Chen, H. I. Lü and Y. N. Yeh, Operations of Interlaced Trees and Graceful Trees, *Southeast Asian Bull. Math.* 21 (1997), 337–348.
- [3] J.A. Gallian, A dynamic survey of graph labeling, *Electron. J. Combin.* DS 6 (2010), 1–296.
- [4] F. Harary, *Graph theory*, Addition Wesley, Academic press, 1972.
- [5] S.M. Hegde, Set colorings of graphs, *European J. Combin.* 30(4) (2009), 986–995.
- [6] M. Mollard and C. Payan, On two conjectures about set graceful graphs, *European J. Combin.* 10(2) (1989), 185–187.
- [7] D.B. West, *Introduction to Graph Theory*, Prentice Hall of India, 2001.

(Received 23 Aug 2010; revised 19 Nov 2011)