

# Maximal outerplanar graphs as chordal graphs, path-neighborhood graphs, and triangle graphs

R.C. LASKAR

*Department of Mathematical Sciences*  
*Clemson University*  
*Clemson, SC 29634*  
*U.S.A.*  
rclsk@clemson.edu

HENRY MARTYN MULDER

*Econometrisch Instituut, Erasmus Universiteit*  
*P.O. Box 1738, 3000 DR Rotterdam*  
*The Netherlands*  
hmulder@ese.eur.nl

B. NOVICK

*Department of Mathematical Sciences*  
*Clemson University*  
*Clemson, SC 29634*  
*U.S.A.*  
nbeth@clemson.edu

## Abstract

Maximal outerplanar graphs are characterized using three different classes of graphs. A path-neighborhood graph is a connected graph in which every neighborhood induces a path. The triangle graph  $T(G)$  has the triangles of the graph  $G$  as its vertices, two of these being adjacent whenever as triangles in  $G$  they share an edge. A graph is edge-triangular if every edge is in at least one triangle. The main results can be summarized as follows: the class of maximal outerplanar graphs is precisely the intersection of any of the two following classes: the chordal graphs, the path-neighborhood graphs, the edge-triangular graphs having a tree as triangle graph.

## 1 Introduction

The problem of characterizing families of graphs is ubiquitous in the literature, and has a long tradition. There is Kuratowski's well-known characterization of planar graphs [17] dating back to 1930. There is the recent breakthrough result, characterizing perfect graphs, due to Chudnovski et al. [7], first announced in 2002 and published in 2006, settling the longstanding conjecture of Berge [3]. A gamut of contributions appeared between these two events; see the survey on graph classes by Brandstädt, Le and Spinrad [5]. In fact, Kuratowski's result was by no means the first. The characterization of Eulerian graphs as connected graphs in which each vertex has even degree has its ultimate origins in Euler's Seven Bridges of Königsberg problem of 1736 [10].<sup>1</sup>

One often very useful way to characterize a given family of graphs is to show that it is precisely the intersection of two or more other well-known classes of graphs. For example, in 1977, Földes and Hammer [12] show that the family of split graphs, namely those graphs whose vertex set may be partitioned into a clique and an independent set, is precisely the intersection of the class of chordal graphs and the class of co-chordal graphs. A motivation here was that split graphs are necessarily perfect graphs and the reader is directed to [5] for many more examples involving perfect graphs.

Trees are a classical example of a class of graphs definable by means of intersection of graph classes. As is well known, the family of trees is precisely the intersection of any of two of the following three classes of graphs: the connected graphs, the acyclic graphs, and the graphs having one edge fewer than vertices. On the other hand, neither connectivity, acyclicity nor the property of having one edge fewer than vertices, suffices for a graph to be a tree. Indeed, any one of these classes contains an infinite number of graphs that are not trees. In the present paper, we offer an analogous characterization for maximal outerplanar graphs.

An outerplanar graph is a planar graph that has a plane embedding such that all vertices lie on the boundary of the outer face. A maximal outerplanar graph is an outerplanar graph such that the number of edges is maximum. Another way to view a maximal outerplanar graph with at least three vertices is that it is the triangulation of a plane cycle. Because of their simple and nice structure, maximal outerplanar graphs, also known as MOP's, have attracted much attention in the literature. Many structural and computational results are available. For a selection of the literature see [21, 2, 15]. This paper involves characterizations of maximal outerplanar graphs in terms of three different classes of graphs. The first class is that of the *chordal graphs*. These are well known and well studied: a graph is chordal if it does not

---

<sup>1</sup>Note that Euler did not use graphs; these were only introduced in 1878 by Sylvester. Moreover, Euler only proved the trivial implication of the if-and-only-if statement. A full proof, in terms of drawings of closed lines in the plane, was given first by Hierholzer in 1873. The first graph theoretic formulation and proof was given by E. Lucas in 1891. Also in 1891 Julius Petersen proved the graph result, probably independently of the tradition of the Königsberg Bridges Problem. See [22] for details.

contain an induced cycle of length at least 4. They were introduced as rigid circuit graphs by Dirac [8], who gave the fundamental characterization that chordal graphs are precisely the graphs admitting a simplicial elimination ordering. The second class is that of the ‘path-neighborhood graphs’, introduced in [18]: a *path-neighborhood graph* is a connected graph in which every neighborhood induces a path. The third class involves triangle graphs, introduced by [24]; see also [9, 1, 19]. Let  $G$  be a graph. The *triangle graph*  $T(G)$  of  $G$  has the triangles of  $G$  as its vertices, and two vertices of  $T(G)$  are adjacent whenever as triangles in  $G$  they share an edge. The third class we have in mind is the class of graphs  $G$  such that every edge of  $G$  is in a triangle and  $T(G)$  is a tree. The main results of this paper can be summarized as follows. The maximal outerplanar graphs form a proper subclass of each of these three classes, but the intersection of any two of these three classes consists precisely of the class of maximal outerplanar graphs.

## 2 Maximal outerplanar graphs and three classes

An *outerplanar graph* is a planar graph that allows an embedding in the plane such that all vertices are on the outer face. In the sequel we will always assume that such a plane embedding is given. A *maximal outerplanar graph* is an outerplanar graph with a maximum number of edges. In the plane embedding the boundary of the outer face, provided it has at least three vertices, is then a hamiltonian cycle. All other edges form a triangulation of this outer cycle. Outerplanar graphs occur for the first time in the literature in Harary’s classical book [14]. The following theorem appeared in [20]. Its proof is an easy exercise: maximality implies that each induced cycle is a triangle. By definition a path is a sequence of alternatingly vertices and edges, where each edge joins its preceding vertex with its succeeding vertex. For readability, we replace edges by arrows.

**Theorem 1** *Let  $G$  be a maximal outerplanar graph with its plane embedding, and let  $v$  be any vertex. Then the neighborhood of  $v$  consists of an induced path  $v_1 \rightarrow v_2 \rightarrow \dots \rightarrow v_k$ , where  $v_i \rightarrow v_{i+1}$  means  $v_i v_{i+1}$  is an edge. The edges  $vv_1$  and  $vv_k$  are on the outer face, whereas the edges  $vv_2, vv_3, \dots, vv_k$  are interior edges.*

Let  $G$  be a graph, and let  $u$  be a vertex of  $G$ . The *neighborhood*  $N(u)$  of  $u$  is the set of neighbors of  $u$ . A *path-neighborhood graph* is a connected graph in which the neighborhood of each vertex induces a path. Path-neighborhood graphs were introduced in [18]. Theorem 1 says that all maximal outerplanar graphs are path-neighborhood graphs.

Let  $\mathcal{P}$  be a property of a vertex in a graph  $G = (V, E)$  of order  $|V| = n$ . A  *$\mathcal{P}$ -elimination ordering* of  $G$  is an ordering  $v_1, v_2, \dots, v_n$  of  $V$  such that  $v_i$  has property  $\mathcal{P}$  in the subgraph of  $G$  induced by  $v_i, v_{i+1}, \dots, v_n$ , for  $i = 1, 2, \dots, n-1$ . For instance, a *simplicial vertex* is a vertex, the neighborhood of which is a clique. Then a *simplicial elimination ordering* of  $G$  is a  $\mathcal{P}$ -elimination ordering in which property  $\mathcal{P}$  is “being simplicial”. Similarly, a *degree-two elimination ordering* of  $G$  is a  $\mathcal{P}$ -elimination

ordering in which property  $\mathcal{P}$  is “having degree 2”. With such a property we have to amend: in the last step of the elimination order  $v_{n-1}$  does not have degree 2 anymore for obvious reasons: there are only two vertices left. So now we require that  $v_{n-1}$  has degree as close to 2 as possible, so  $v_{n-1}v_n$  is an edge. Note that a graph admitting a degree-two elimination ordering necessarily is connected.

A *chordal graph* is a graph without induced cycles of length at least 4. In 1961 Dirac [8] proved the classical result that a graph is chordal if and only if it admits a simplicial elimination ordering. For more information on chordal graphs see [13].

As observed above, each induced cycle in a maximal outerplanar graph is a triangle. This implies the well-known fact that a maximal outerplanar graph is chordal. Furthermore, any simplicial vertex having degree 3 or more will be part of a  $K_4$  subgraph, a well-known obstruction for outerplanar graphs, see [6]. In light of these considerations, we can formulate another result, which is part of folklore.

**Theorem 2** *A maximal outerplanar graph is necessarily chordal, admitting a simplicial elimination ordering which is at the same time a degree-two elimination ordering.*

Let us call an elimination ordering that is both simplicial and degree-two a *triangle elimination ordering*. This seems appropriate because by eliminating such a vertex we destroy a triangle by deleting a vertex from the triangle that has no neighbors outside the triangle. Closely related is the concept of a *2-tree*, see [23], namely, a graph constructed by beginning with  $K_2$  and at each iteration adding a new vertex  $v$ , joining it to two existing, adjacent vertices, thereby forming a new triangle. While it is not essential to our main results, we next note, in passing, a strengthening of the above result. In essence, this result is proved earlier, from an algorithmic point of view, and within the proof of a different result, see [23]. We offer our version of the proof for clarity and brevity.

**Theorem 3** *A graph is maximal outerplanar if and only if it admits a triangle elimination ordering and does not contain a  $K_{1,1,3}$ .*

**Proof.** Let  $G$  be a maximal outerplanar graph. Then obviously  $G$  does not contain a  $K_{1,1,3}$ , since the obstruction  $K_{2,3}$  is a subgraph of  $K_{1,1,3}$ . The existence of the triangle elimination ordering follows from Theorem 2.

Conversely, let  $G$  be  $K_{1,1,3}$ -free having a triangle elimination scheme. We proceed by induction on the number  $n$  of vertices. For  $n \leq 3$  it is obvious that  $G$  is maximal outerplanar. So let  $n \geq 4$ , and let  $v$  be a simplicial vertex of degree 2 in  $G$ . Let  $x$  and  $y$  be the neighbors of  $v$ , so that  $xy$  is an edge. By induction  $G - v$  is maximal outerplanar. It suffices to prove that  $xy$  is on the outer-cycle in a plane embedding of  $G - v$ . Assume to the contrary that  $xy$  is a chord of the outer-cycle. Then we can find vertices  $p$  and  $q$  such that  $x, y, p$  and  $x, y, q$  induce triangles on different sides of  $xy$  in the plane embedding of  $G - v$ . But now  $v, x, y, p, q$  induce a  $K_{1,1,3}$  in  $G$ . This impossibility completes the proof.  $\square$

As observed, a maximal outerplanar graph is a chordal graph as well as a path-neighborhood graph. The following result states that together these properties suffice. A related (but not identical) result, with a somewhat longer proof, is that  $G$  is maximal outerplanar if and only if  $G$  is a path-neighborhood graph that is ‘2-degenerate,’ namely, every subgraph of  $G$  has a vertex of degree 2 or less, see [16].

**Theorem 4** *A graph  $G$  is a chordal path-neighborhood graph if and only if  $G$  is a maximal outerplanar graph.*

**Proof.** Theorems 1 and 2 tell us that a maximal outerplanar graph is a chordal path-neighborhood graph.

We prove the converse by induction on the number of vertices  $n$ . For  $n \leq 3$ , the theorem is trivial. So assume that  $n \geq 4$ . Let  $v$  be the first vertex in a simplicial elimination ordering, making  $v$  simplicial and  $G - v$  still chordal. Since the neighborhood of  $v$  is both an induced path and an induced clique, it must be an edge  $xy$ . Then  $N(x)$  induces the path  $P_x = v \rightarrow y \rightarrow \dots$ , and  $N(y)$  induces the path  $P_y = v \rightarrow x \rightarrow \dots$ . So in  $G - v$  the neighborhood of  $x$  induces the path  $P_x - v = y \rightarrow \dots$ , and the neighborhood of  $y$  induces the path  $P_y - v = x \rightarrow \dots$ . Hence  $G - v$  is again a chordal, path-neighborhood graph, so that, by induction,  $G - v$  is a maximal outerplanar graph. By Theorem 1, the edge  $xy$  is on its outer face. This implies that  $G$  is also maximal outerplanar.  $\square$

The *triangle graph*  $T(G)$  of a graph  $G$  is the graph with the triangles of  $G$  as vertices, and two such vertices are joined in  $T(G)$  if, as triangles in  $G$ , they share an edge. Triangle graphs were first introduced in a different context by Pullman [24]. They were introduced later independently a couple of times, see e.g. [9, 1, 19]. The following fact follows easily from the definition of  $T(G)$ .

**Fact 5**  $K_{1,4}$  does not occur in  $T(G)$  as an induced subgraph.

Let  $G$  be a graph, and let  $u$  be a vertex of  $G$ . Assume that  $N(u)$  induces a disconnected graph, consisting say of two disjoint subgraphs  $N_1$  and  $N_2$  with no edges joining  $N_1$  and  $N_2$ . Let  $H$  be the graph obtained from  $G$  by replacing  $u$  in  $G$  by two new vertices  $u_1, u_2$  and joining  $u_i$  to all vertices in  $N_i$ , for  $i = 1, 2$ . Note that in  $H$  the distance between  $u_1$  and  $u_2$  is at least 4. We say that  $H$  is obtained from  $G$  by *splitting*  $u$ . Clearly, we have  $T(G) \cong T(H)$ . By successive splittings we can get a graph  $\widehat{G}$  from  $G$  in which all neighborhoods are connected. In fact, as we show in the next result,  $\widehat{G}$  is independent of the order of the splittings and is hence unique. Clearly we have  $T(G) \cong T(\widehat{G})$ .

**Proposition 6** *The splitting operation on any graph is order independent.*

**Proof.** It suffices to show that any splitting operation preserves the connected components of neighborhoods for all vertices.

Assume to the contrary that in splitting a vertex  $u$ , replacing it with  $u_1$  and  $u_2$ , some other vertex  $w$  has neighbors  $x$  and  $y$  which were in the same connected component of  $N(w)$  prior to the split, but afterwards are in different ones. But then, prior to the split,  $x$  was connected to  $y$  via a path in  $N(w)$ . This path was destroyed in splitting  $u$ . So this path must have contained  $u$ . No edges are destroyed in splitting  $u$ . So  $u_1w$  and  $u_2w$  both are edges after  $u$  is split, which is impossible.  $\square$

From the viewpoint of constructing the triangle graph  $T(G)$  of a graph  $G$ , any vertex or edge in  $G$  not contained in a triangle is irrelevant, and may be deleted. Therefore we restrict ourselves to graphs in which every edge is contained in a triangle and that have no isolates. We call such a graph *edge-triangular*, for want of a better term. Note that a maximal outerplanar graph with at least three vertices is necessarily an edge-triangular graph.

By Proposition 6, for any edge-triangular graph  $G$  it makes sense to define  $\widehat{G}$  to be the unique graph obtained from  $G$  by successive splittings and having all its neighborhoods connected. On the class of connected edge-triangular graphs we may also define the relation  $\sim$  by  $G \sim H$  if  $\widehat{H} \cong \widehat{G}$ . The relation  $\sim$  is clearly an equivalence relation. Note that two equivalent graphs have the same triangle graph, and any equivalence class contains a unique graph with connected neighborhoods, viz.  $\widehat{G}$ , for any graph  $G$  in the class.

Let  $G$  be a maximal outerplanar graph having at least three vertices. By Fact 5, it is easy to see that  $T(G)$  is a tree of maximum degree at most 3. It can also be obtained from the dual graph of  $G$  by deleting the vertex that represents the outer face of  $G$ . This graph is the so-called *weak dual* of  $G$ , see [11]. It can be obtained from  $G$  as follows: the interior faces of  $G$  are the vertices of the weak dual  $G^*$ , two vertices in  $G^*$  being adjacent whenever as faces in  $G$  they share an edge in their boundaries. The weak dual was used in [2] to construct recognition algorithms for outerplanar graphs. And it was used in [15] to study maximal outerplanar graphs and their interior graphs: the graph obtained by deleting the edges on the exterior face of the maximal outerplanar graph.

**Theorem 7** *Let  $G$  be an edge-triangular graph. Then  $T(G)$  is a tree if and only if  $\widehat{G}$  is a maximal outerplanar graph.*

**Proof.** As observed above, if  $\widehat{G}$  is a maximal outerplanar graph, then  $T(G) \cong T(\widehat{G})$  is a tree.

Conversely, let  $G$  be an edge-triangular graph such that  $T(G)$  is a tree. Recall that  $T(G) \cong T(\widehat{G})$ . In  $\widehat{G}$  all neighborhoods are connected. We prove by induction on the number of vertices  $n$  of  $T(\widehat{G})$  that  $\widehat{G}$  is a maximal outerplanar graph. First note that, by Fact 5, the maximum degree in  $T(\widehat{G})$  is at most 3.

If  $n = 1$ , then  $\widehat{G}$  is a triangle, and we are done. So assume that  $n \geq 2$ . Then  $T(\widehat{G})$  contains a pendant vertex  $x$  adjacent to a vertex  $y$ . Let  $x$  represent the triangle

in  $\widehat{G}$  on  $a, b, c$ , and let  $y$  represent the triangle in  $\widehat{G}$  on  $b, c, d$ . Then the edges  $ab$  and  $ac$  in  $\widehat{G}$  are not contained in any other triangle, so  $a$  is a vertex of degree 2 in  $\widehat{G}$ . Moreover, edge  $bc$  is contained only in the triangles representing  $x$  and  $y$ . So in  $\widehat{G} - a$  edge  $bc$  is contained in a unique triangle. Now  $T(\widehat{G}) - x$  is the triangle graph of  $\widehat{G} - a$ . Moreover,  $\widehat{G}$  being an edge-triangular graph with connected neighborhoods, it follows that  $\widehat{G} - a$  is again such. So, by induction,  $\widehat{G} - a$  is a maximal outerplanar graph. Since edge  $bc$  is in a unique triangle in  $\widehat{G} - a$ , it must be on the outer face of  $\widehat{G} - a$ . Hence, if we add  $a$  back on, then  $\widehat{G}$  remains a maximal outerplanar graph.  $\square$

A *snake* is a maximal outerplanar graph in which every triangle shares an edge with the outer face. Clearly, its triangle graph is a path.

**Corollary 8** *Let  $G$  be a edge-triangular graph. Then  $T(G)$  is a path if and only if  $\widehat{G}$  is a snake.*

The following new characterization of maximal outerplanar graphs is an easy consequence of Theorem 7.

**Theorem 9** *An edge-triangular graph  $G$  is a path-neighborhood graph with a tree as its triangle graph if and only if  $G$  is a maximal outerplanar graph.*

**Proof.** If  $G$  is a path-neighborhood graph, then  $\widehat{G} = G$ , and Theorem 7 tells us that,  $T(G)$  being a tree,  $G$  is maximal outerplanar.

The converse follows from Theorems 7 and 1.  $\square$

Finally, we consider the intersection of the class of chordal graphs and that of the graphs with a tree as triangle graph.

**Theorem 10** *An edge-triangular graph  $G$  is a chordal graph with a tree as its triangle graph if and only if  $G$  is a maximal outerplanar graph.*

**Proof.** Let  $G$  be a maximal outerplanar graph. Then it follows from Theorems 4 and 7 that  $G$  is chordal and has a tree as its triangle graph.

Conversely, let  $G$  be a chordal graph with  $T(G)$  a tree. Then,  $T(G)$  being a tree,  $K_4$  and  $K_{1,1,3}$  do not occur in  $G$ . Furthermore a simplicial vertex in  $G$  is of degree 2, and the triangle containing the simplicial vertex is a pendant vertex in  $T(G)$ . We use induction on the number  $n$  of vertices of  $G$ . For  $n \leq 3$  the assertion is trivial. So assume that  $n \geq 4$ , and let  $v$  be a simplicial vertex of  $G$  with neighbors  $x$  and  $y$ . Then  $v, x, y$  form a triangle represented by a pendant vertex  $p$  in  $T(G)$ . Let its neighbor  $q$  in  $T(G)$  represent the triangle in  $G$  on  $x, y, z$ . Now  $G - v$  is a chordal graph with  $T(G) - p$  as its triangle graph. So, by induction,  $G - v$  is maximal outerplanar. If edge  $xy$  is on the outer face of  $G - v$ , then  $G$  is maximal outerplanar

as well. Consider the neighborhood  $N(x)$  of  $x$  in  $G - v$ . Since  $G - v$  is maximal outerplanar,  $N(x)$  induces a path  $P_x$ . If  $y$  would be internal vertex of this path, say, with neighbors  $z$  and  $w$ , then  $v, x, y, z, w$  would produce a  $K_{1,1,3}$  in  $G$ , contradicting Theorem 7. So  $y$  is not an internal vertex of  $P_x$ . So, by Theorem 1, the edge  $xy$  is on the outer face, and we are done.  $\square$

### 3 Concluding Remarks




Figure 1: Maximal outerplanar graphs and the three classes

In this paper we have considered three classes of graphs, namely, chordal graphs, path-neighborhood graphs, and edge-triangular graphs having a tree as triangle graph. Note that, by definition, a path-neighborhood graph is connected. Moreover, if a graph  $G$  is edge-triangular, then its triangle graph being connected implies that  $G$  itself is also connected. In the previous sections we have proved that the intersection of any two of these classes constitutes precisely the class of maximal outerplanar graphs. This is depicted in Figure 1. As part of our concluding remarks, we present examples that show that the class of maximal outerplanar graphs is properly contained in each of the three classes. In fact, each of the three classes contains an infinite number of graphs that are not maximal outerplanar.

Any complete graph with more than three vertices is chordal but not a path-neighborhood graph and its triangle graph contains a  $K_4$ .

The triangulated band  $Z_n$  in [18] is a path-neighborhood graph, but it is not chordal, and its triangle graph is the  $n$ -cycle. For a picture of  $Z_4$  see Figure 2.

Take a snake, in which the vertices of degree 2 are at distance at least 4. Glue the two vertices of degree 2 together. This is not a chordal graph and not a path-neighborhood graph, but its triangle graph is a path. See Figure 2 for an example.






Figure 2: Two non-outerplanar graphs

Lastly, we discuss some open problems. There is a rich and still growing literature on chordal graphs. So far not much is known about the other two classes: the path-neighborhood graphs and the triangle graphs. It seems there are still many interesting open problems involving these classes and related issues. We mention a few of these problems here.

### Problem 1

It is well-known that a graph is chordal if and only if it admits a simplicial elimination ordering. As we have seen in the current paper, there are many other, interesting, elimination orderings. For example, as we showed in Theorem 3, a graph is maximal outerplanar if and only if it admits a triangle elimination ordering and does not contain a  $K_{1,1,3}$ . Is there a nice characterization of graphs admitting a degree two elimination ordering? Note that a triangle elimination ordering is a degree two elimination ordering.

### Problem 2

Boros, Jamison, Laskar and Mulder [4] show the existence of so-called 3-simplicial vertices in planar graphs: a vertex is 3-simplicial if its neighborhood can be edge-covered by at most three cliques. Here the analogue for outerplanar graphs is a simple one: an outerplanar graph always contains a vertex of degree at most two, which is 2-simplicial if it has two non-adjacent neighbors; otherwise it is simplicial. It may be fruitful to study, more generally, 2-simplicial orderings and graphs which admit 2-simplicial orderings.

### Problem 3

In Theorem 9 we prove that an edge-triangular graph  $G$  is maximal outerplanar if and only if  $G$  is a path-neighborhood graph with a tree as its triangle graph. It is likely that more could be said about the triangle graph of a path-neighborhood graph. In other words, suppose  $G$  is a path-neighborhood graph with triangle graph  $T(G)$ . By Theorem 9,  $G$  is maximal outerplanar if and only if  $T(G)$  is a tree. Question: Characterize those graphs where, say, (a)  $T(G)$  is a cycle; (b)  $T(G)$  is connected; and so forth.

### References

- [1] R. Balakrishnan, *Triangle graphs*, in: Graph connections (Cochin, 1998), p. 44, Allied Publ., New Delhi, 1999.
- [2] T. Beyer, W. Jones and S.L. Mitchell, Linear algorithms for isomorphisms of maximal outerplanar graphs, *J. Assoc. Computing Machinery* **26** (1979), 603–610.
- [3] C. Berge (1961). *Färbung von Graphen, deren sämtliche bzw. deren ungerade Kreise starr sind*, *Wiss. Z. Martin-Luther-Univ. Halle-Wittenberg Math.-Natur. Reihe* **10** (1961), 114.
- [4] E. Boros, R.E. Jamison, R.C. Laska and H.M. Mulder, On 3-simplicial vertices in planar graphs, *Discuss. Math. Graph Theory* **14** (2004), 413–421.
- [5] A. Brandstädt, V.B. Le and J.P. Spinrad, *Graph Classes: a survey*, SIAM Monographs on Discrete Mathematics and Applications. Philadelphia, 1999.
- [6] G. Chartrand and F. Harary, Planar permutation graphs, *Ann. Inst. Henri Poincaré B* **3** (1967), 433–438.
- [7] M. Chudnovsky, N. Robertson, P.D. Seymour and R. Thomas, The strong perfect graph theorem, *Ann. Math.* **164** (2006), 51–229.
- [8] G.A. Dirac, On rigid circuit graphs, *Abh. Math. Sem. Univ. Hamburg* **25** (1961), 71–76.
- [9] Y. Egawa and R.E. Ramos, Triangle graphs, *Math. Japon.* **36** (1991), 465–467.
- [10] L. Euler, Solutio problematis ad geometriam situs pertinentis, *Commentarii Academiae Scientiarum Imperialis Petropolitanae* **8** (1736), 128–140.
- [11] H.J. Fleischner, D.P. Geller and F. Harary, Outerplanar graphs and weak duals, *J. Indian Math. Soc.* **38** (1974), 215–219.
- [12] S. Földes and P.L. Hammer, Split graphs, *Congressus Numer.* **19** (1977), 311–315.

- [13] M.C. Golumbic, *Algorithmic Graph Theory and Perfect Graphs*, Second ed., Annals of Discrete Math. **57**, Elsevier, Amsterdam, 2004.
- [14] F. Harary, *Graph Theory*, Addison-Wesley, Reading Massachusetts, 1969.
- [15] S.M. Hedetniemi, A. Proskurowski and M.M. Sysło, Interior graphs of maximal outerplane graphs, *J. Combin. Theory Ser. B* **38** (1985), 156–167.
- [16] G. Hopkins and W. Staton, Outerplanarity without topology, *Bull. Inst. Combin. Appl.* **21** (1997), 112–116.
- [17] K. Kuratowski, Sur le problème des courbes gauches en topologie, *Fund. Math.* **15** (1930), 271–283.
- [18] R.C. Laskar and H.M. Mulder, Path-neighborhood graphs, preprint, *EI-Report EI 2010-64*, Econometrisch Instituut, Erasmus Universiteit, Rotterdam.
- [19] R.C. Laskar, H.M. Mulder, Triangle graphs, *EI-Report EI 2009-42*, Econometrisch Instituut, Erasmus Universiteit, Rotterdam.
- [20] S.L. Mitchell, *Algorithms on trees and maximal outerplanar graphs: design, complexity analysis and data structures study*, PhD thesis, University of Virginia, 1977.
- [21] S.L. Mitchell, Linear algorithms to recognize outerplanar and maximal outerplanar graphs, *Inform. Proc. Letters* **9** (1979), 229–232.
- [22] H.M. Mulder, *Die Entsehung der Graphentheorie*, in: K. Wagner, R. Bodendiek, Graphentheorie, Band 3, Wissenschaftsverlag, Mannheim, 1992, pp. 296–313, 381–383.
- [23] A. Proskurowski, Minimum dominating cycles in 2-trees, *Internat. J. Comput.* **8** (1979), 405–417.
- [24] N. Pullman, Clique covering of graphs. IV. Algorithms, *SIAM J. Comput.* **13** (1984), 57–75.

(Received 5 May 2011; revised 12 Oct 2011)