

Kostka multiplicity one for multipartitions

JAMES JANOPAUL-NAYLOR C. RYAN VINROOT

*Department of Mathematics
College of William and Mary
P. O. Box 8795, Williamsburg, VA 23187
U.S.A.*

jrjanopaulnayl@email.wm.edu vinroot@math.wm.edu

Abstract

If $[\lambda(j)]$ is a multipartition of the positive integer n (a sequence of partitions with total size n), and μ is a partition of n , we study the number $K_{[\lambda(j)]\mu}$ of sequences of semistandard Young tableaux of shape $[\lambda(j)]$ and total weight μ . We show that the numbers $K_{[\lambda(j)]\mu}$ occur naturally as the multiplicities in certain permutation representations of wreath products. The main result is a set of conditions on $[\lambda(j)]$ and μ which are equivalent to $K_{[\lambda(j)]\mu} = 1$, generalizing a theorem of Berenshteĭn and Zelevinskiĭ. Finally, we discuss some computational aspects of the problem, and we give an application to multiplicities in the degenerate Gel'fand-Graev representations of the finite general linear group. We show that the problem of determining whether a given irreducible representation of the finite general linear group appears with nonzero multiplicity in a given degenerate Gel'fand-Graev representation, with their partition parameters as input, is NP -complete.

1 Introduction

Young tableaux and Kostka numbers, in their various forms, are of central importance in combinatorics and representation theory [21]. In symmetric function theory, Kostka numbers appear in transition matrices between fundamental bases, and in representation theory they appear as multiplicity coefficients in numerous contexts. An important question in both of these settings is whether a basic object appears with *multiplicity one* in an expansion, since this uniqueness can be exploited to identify interesting properties of this object. It is with this motivation that A. D. Berenshteĭn and A. V. Zelevinskiĭ [2] solve the following problem: given a complex semisimple Lie algebra \mathfrak{g} and an irreducible \mathfrak{g} -module V_λ with highest weight λ , find all weights ω such that the weight subspace $V_\lambda(\omega)$ of V_λ has dimension 1. In the case that \mathfrak{g} is of type A , the dimension of the weight subspace $V_\lambda(\omega)$ is exactly the Kostka number $K_{\lambda\omega}$, or the number of semistandard Young tableaux of shape λ and content ω (which we will call the *weight* of the tableau henceforth, and we will not refer to

weights in the Lie algebra context again). As an important corollary, Berenshteĭn and Zelevinskiĭ thus obtain conditions on partitions λ and μ of a positive integer n which are equivalent to the statement that $K_{\lambda\mu} = 1$. The main purpose of this paper is to generalize this result in the case that we replace λ with a multipartition $[\lambda(j)]$, that is, a sequence of partitions with total size n , which we accomplish in Theorem 5.3.

The organization and results of this paper are as follows. In Section 2, we give background material and the main definitions, starting with partitions and tableaux (Section 2.1), Kostka numbers and some classical results on symmetric functions and the symmetric group (Section 2.2), and the definitions for multipartitions and multitableaux, including the basic results which parallel those for classical Kostka numbers (Section 2.3). In Section 3, we state the relevant result of Berenshteĭn and Zelevinskiĭ for Kostka numbers associated with partitions in Theorem 3.1. In an effort to keep this paper as self-contained as possible, we give a complete proof of Theorem 3.1 using tableau combinatorics, as there does not seem to be a tableau-theoretic proof of this statement in the literature previously.

In Section 4, we turn to the representation theory of wreath products of finite groups to find motivation to study Kostka numbers associated with multipartitions. The main result here is Theorem 4.5, where we decompose wreath product permutation characters which are analogous to the symmetric group on a Young subgroup, and we find that multipartition Kostka numbers appear as multiplicities. In Section 5, we prove Theorem 5.3, in which we give conditions on a multipartition $[\lambda(j)]$ and a partition μ which are equivalent to the statement $K_{[\lambda(j)]\mu} = 1$. In Corollary 5.4, we classify exactly which multipartitions $[\lambda(j)]$ are such that there exists a unique partition μ satisfying $K_{[\lambda(j)]\mu} = 1$, generalizing the result for partitions given in Corollary 3.4.

Lastly, in Section 6, we consider some computational problems related to the main results. In particular, it is known from more general results that the question of whether the Kostka number $K_{\lambda\mu}$ is equal to any particular value can be answered in polynomial time, and some of these results immediately extend to the multipartition case. We then consider an application to the characters of the finite general linear group by applying our main result Theorem 5.3 to obtain some information on the decompositions of the degenerate Gel'fand-Graev characters. In considering the more general version of Kostka numbers which appear in this application, we show in Proposition 6.3 that the question of whether an irreducible character appears as a constituent of a degenerate Gel'fand-Graev character, with the partition data as input, is an NP -complete problem.

Another generalization of the theorem of Berenshteĭn and Zelevinskiĭ which has been studied is the question of when the *Littlewood-Richardson coefficient* takes the value 1 (see [15, Corollary, Sec. 6]). For partitions λ, μ , and ν , such that $|\nu| + |\mu| = |\lambda|$, the Littlewood-Richardson coefficient $c_{\nu,\mu}^\lambda$ is the number of semistandard Young tableaux of weight μ and skew shape $\lambda - \nu$ whose filling satisfies the *lattice permutation property*, or the *Yamanouchi condition* (see [5, Section 5.1] or [16, Section I.9]). In particular, $K_{\lambda\mu} = c_{\nu,\mu}^\lambda = 1$ when ν is the empty partition and $\lambda = \mu$. More generally, Kostka numbers can be viewed as special kinds of Littlewood-


Richardson coefficients (see [13] for a precise construction). Littlewood-Richardson coefficients are studied and interpreted through the *honeycomb model* by Knutson and Tao [14] and Knutson, Tao, and Woodward [15], with several important combinatorial and representation-theoretic applications. There is some indication [14, After Corollary, Sec. 6] that the results on the honeycomb model in the multipartition case can be carried out using tools of Zelevinsky [25]. It would be interesting to carry out this construction in detail, and understand the main result of this paper in the context of the honeycomb model.

2 Preliminaries


2.1 Partitions and Tableaux

For any non-negative integer n , a *partition* of n is a finite sequence $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_l)$ of positive integers such that $\lambda_i \geq \lambda_{i+1}$ for $i < l$ and $\sum_{i=1}^l \lambda_i = n$, where λ_i is the i th *part* of λ . The number of parts of λ is the *length* of λ , written $\ell(\lambda)$. If λ is a partition of n , we write $\lambda \vdash n$, and we say λ has *size* n , written $|\lambda| = n$. If $n = 0$, the only partition of n is the *empty partition*, written (0) . We let \mathcal{P}_n denote the set of all partitions of size n , and \mathcal{P} the set of all partitions.

We may represent a partition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_l)$ by its *Young diagram*, which has a row of λ_i boxes for each part of λ , which are upper-left justified. Notationally, we will often identify a partition λ with its Young diagram. For example, if $\lambda = (4, 3, 2, 2, 1)$, then $\lambda \vdash 12$ (so $|\lambda| = 12$), $\ell(\lambda) = 5$, and the Young diagram for λ is




Given $\lambda, \mu \in \mathcal{P}$, we say $\mu \subset \lambda$ if $\mu_i \leq \lambda_i$ for all i . That is, $\mu \subset \lambda$ exactly when the Young diagram for μ fits inside of the Young diagram for λ . If $\mu \subset \lambda$, we define $\lambda - \mu$ to be the result of removing the boxes of the Young diagram for μ from the Young diagram of λ , resulting in a *skew diagram*. For example, if $\lambda = (5, 4, 4, 1)$ and $\mu = (3, 2, 1)$, then $\mu \subset \lambda$, and the Young diagram for λ with the Young diagram for μ inside of it marked by dots, and the skew diagram for $\lambda - \mu$, are




The size of a skew diagram is defined to be the number of boxes it has, and above we have $|\lambda - \mu| = 8$. A *horizontal strip* is a skew diagram which has at most one box in each column, and a horizontal m -strip is a horizontal strip of size m . For example,

if $\lambda = (5, 4, 1)$ and $\mu = (4, 2)$, then $\lambda - \mu$ is a horizontal 4-strip with diagram



There is a partial order on the set \mathcal{P}_n called the *dominance partial order* defined as follows. Given two partitions $\lambda, \mu \in \mathcal{P}_n$, define $\lambda \trianglerighteq \mu$ if for each $k \geq 1$, $\sum_{i \leq k} \lambda_i \geq \sum_{i \leq k} \mu_i$. For example, if $\lambda = (3, 2, 1)$, $\mu = (3, 1, 1, 1)$, and $\nu = (2, 2, 2)$, then $\lambda \trianglerighteq \mu$ and $\lambda \trianglerighteq \nu$, but μ and ν are incomparable in the dominance partial order.

Given $\lambda \in \mathcal{P}$, a *semistandard Young* (or *column-strict*) *tableau* of *shape* λ is a filling of the boxes of the Young diagram for λ with positive integers, such that entries in each row weakly increase from left to right and entries in each column strictly increase from top to bottom. If T is a semistandard Young tableau of shape λ , the *weight* of T , written $\text{wt}(T)$, is the finite sequence $\omega = (\omega_1, \omega_2, \dots, \omega_l)$, where ω_i is the number of times i is an entry in T , and l is the largest entry in T . Note that $\sum_{i=1}^l \omega_i = |\lambda|$, so if $\lambda \vdash n$, then $\text{wt}(T)$ is a *composition* of n (where we allow entries to be 0, and the length l is the position of the last positive part). We will mainly concern ourselves with the case that the weight of a tableau is also a partition, so that $\omega_i \geq \omega_{i+1}$ for each i , and we will typically denote the weight by μ in this case. An example of a semistandard Young tableau of shape $\lambda = (4, 2, 2, 1)$ and weight $\mu = (3, 3, 2, 1)$ is



We may also describe a semistandard Young tableau of shape λ and weight ω to be a sequence

$$(0) = \lambda^{(0)} \subset \lambda^{(1)} \subset \dots \subset \lambda^{(l)} = \lambda$$

of nested partitions, such that each $\lambda^{(i)} - \lambda^{(i-1)}$ is a (possibly empty) horizontal strip, and $\omega_i = |\lambda^{(i)} - \lambda^{(i-1)}|$. The idea is that the horizontal strip $\lambda^{(i)} - \lambda^{(i-1)}$ is exactly the set of boxes with entry $i > 0$ in the corresponding semistandard Young tableau, and we take $\lambda^{(i)} = \lambda^{(i-1)}$ when there are no i entries.

2.2 Kostka numbers

Let $\lambda \in \mathcal{P}_n$ and let $\omega = (\omega_1, \omega_2, \dots, \omega_l)$ be a composition of n . The number of semistandard Young tableau of shape λ and weight ω is the *Kostka number*, denoted $K_{\lambda\omega}$. Kostka numbers play an important role in algebraic combinatorics and representation theory. A crucial example is in symmetric function theory (see [16, I.2] for an introduction). For $m \geq 0$, let h_m be the complete symmetric function, and if $\mu, \lambda \vdash n$, where $\mu = (\mu_1, \mu_2, \dots, \mu_l)$, let $h_\mu = h_{\mu_1} \cdots h_{\mu_l}$ and let s_λ denote the Schur symmetric function corresponding to λ . Then the Kostka numbers make up the entries of the transition matrix between these two bases of symmetric functions, in that we have

$$h_\mu = \sum_{\lambda \vdash n} K_{\lambda\mu} s_\lambda. \tag{2.1}$$

immediate that for $k < j$, we have $\sum_{i=1}^k \lambda_i^* \geq \sum_{i=1}^k \mu_i^*$, since $\lambda \triangleright \mu$. In the case that $\mu_l \leq \lambda_l$, then $j = l$ and $r_l = \mu_l$, and $\lambda^* \triangleright \mu^*$ follows immediately. So we assume now that $\mu_l - \lambda_l = d > 0$. Consider k such that $j \leq k < l$, and we must show $\sum_{i=1}^k \lambda_i^* \geq \sum_{i=1}^k \mu_i^*$, where $\mu_i^* = \mu_i$ when $i \leq k < l$. Note that whenever $k < i < l$, then $\lambda_i < \mu_l$ since $i > j$, and $\mu_i \leq \mu_l$ since μ is a partition. Thus $\lambda_i - \mu_i \leq 0$, while $\lambda_l - \mu_l = -d$, so that $\sum_{i=k+1}^l \lambda_i - \sum_{i=k+1}^l \mu_i \leq -d$. Since $\sum_{i=1}^l \lambda_i - \sum_{i=1}^l \mu_i = 0$, then we obtain

$$\sum_{i=1}^k \lambda_i - \sum_{i=1}^k \mu_i \geq d. \tag{2.2}$$

Since $\lambda_l < \mu_l$, then we have $r_l = \lambda_l$, and so $\mu_l - r_l = \sum_{i=j}^{l-1} r_i = \mu_l - \lambda_l = d$. So $\sum_{i=j}^k r_i \leq d$. From this fact and (2.2), we now have

$$\sum_{i=1}^k \lambda_i^* - \sum_{i=1}^k \mu_i^* = \sum_{i=1}^k \lambda_i - \sum_{i=1}^k \mu_i - \sum_{i=j}^k r_i \geq d - d = 0.$$

Thus $\lambda^* \triangleright \mu^*$ as claimed, and we can construct a semistandard Young tableau of shape λ and weight μ in this way whenever $\lambda \triangleright \mu$.

Now consider the symmetric group S_n of permutations of $\{1, 2, \dots, n\}$, and the irreducible complex representations of S_n . These representations are parameterized by \mathcal{P}_n , and we denote the irreducible representation of S_n parameterized by $\lambda \vdash n$ by π^λ , and its character by χ^λ . We adapt the convention of [6, Lecture 4] and [16, I.8] that $\chi^{(n)} = \mathbf{1}$, the trivial character of S_n , and $\chi^{(1,1,\dots,1)}$ is the sign character of S_n . Given $\mu = (\mu_1, \dots, \mu_l) \vdash n$, let S_μ be the *Young subgroup*

$$S_\mu = S_{\mu_1} \times S_{\mu_2} \times \dots \times S_{\mu_l},$$

where S_μ is embedded in S_n such that S_{μ_1} permutes 1 through μ_1 , S_{μ_2} permutes $\mu_1 + 1$ through μ_2 , and so on. If we consider the permutation representation of S_n on S_μ , we obtain the decomposition

$$\text{Ind}_{S_\mu}^{S_n}(\mathbf{1}) = \bigoplus_{\lambda \triangleright \mu} K_{\lambda\mu} \pi^\lambda, \tag{2.3}$$

known as Young’s rule, see [6, Corollary 4.39] for example. For any partition μ , one may see that $K_{\mu\mu} = 1$, so that π^μ appears with multiplicity 1 in the decomposition (2.3). Two natural question arise. Which other π^λ occur with multiplicity 1 in the representation $\text{Ind}_{S_\mu}^{S_n}(\mathbf{1})$, and when is it the case that π^λ appears with multiplicity 1 in $\text{Ind}_{S_\mu}^{S_n}(\mathbf{1})$ only for $\mu = \lambda$? We give answers to these two questions in Section 3.

2.3 Multipartitions and multitableaux

If $n \geq 0$ and $r \geq 1$ are integers, then an r -*multipartition* of n is a sequence of r partitions such that the sum of the sizes of the r partitions is n . We denote an r -multipartition of n by $[\lambda(j)] = [\lambda(1), \lambda(2), \dots, \lambda(r)]$, where $[\lambda(j)]$ has size $|\lambda(j)| =$

$\sum_{j=1}^r |\lambda(j)| = n$. Let $\mathcal{P}_n[r]$ denote the set of all r -multipartitions of n , and let $\mathcal{P}[r]$ denote the set of all r -multipartitions. Given $[\lambda(j)] \in \mathcal{P}_n[r]$, we define the partition

$$\tilde{\lambda} \in \mathcal{P}_n \quad \text{which has parts} \quad \tilde{\lambda}_i = \sum_{j=1}^r \lambda(j)_i, \tag{2.4}$$

so that $\ell(\tilde{\lambda}) = \max\{\ell(\lambda(j)) \mid 1 \leq j \leq r\}$. One may also view $\tilde{\lambda}$ as the partition obtained by taking all columns of all $\lambda(j)$, and arranging them from longest to smallest. For example, suppose $[\lambda(j)] \in \mathcal{P}_{12}[3]$, with $\lambda(1) = (2, 1, 1)$, $\lambda(2) = (2, 2)$, $\lambda(3) = (4)$. Then $\tilde{\lambda} = (8, 3, 1)$, with Young diagrams

$$\lambda(1) = \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \\ \hline \square & \\ \hline \end{array}, \quad \lambda(2) = \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \quad \lambda(3) = \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \end{array}, \quad \text{and} \quad \tilde{\lambda} = \begin{array}{|c|c|c|c|c|c|c|c|} \hline \square & \square & \square & \square & \square & \square & \square & \square \\ \hline \square & \square & \square & & & & & \\ \hline \square & & & & & & & \\ \hline \end{array}.$$

Given $[\lambda(j)] \in \mathcal{P}_n[r]$ and a composition ω of n , a *semistandard Young r -multitableau of shape $[\lambda(j)]$ and weight ω* is a sequence $[T(j)]$ of r semistandard Young tableaux, where each $T(j)$ has shape $\lambda(j)$, and if $\omega(j)$ is the weight of $\lambda(j)$, then ω has parts given by $\omega_i = \sum_{j=1}^r \omega(j)_i$. As was the case with single tableau, we will primarily deal with r -multitableau which have total weight given by a partition μ .

For example, if $[\lambda(j)] \in \mathcal{P}_{12}[3]$ is the same as in the previous example, one semistandard Young 3-multitableau of weight $\mu = (4, 3, 3, 1, 1)$ is given by

$$T(1) = \begin{array}{|c|c|} \hline \boxed{1} & \boxed{3} \\ \hline \boxed{3} & \\ \hline \boxed{4} & \\ \hline \end{array}, \quad T(2) = \begin{array}{|c|c|} \hline \boxed{1} & \boxed{2} \\ \hline \boxed{2} & \boxed{3} \\ \hline \end{array}, \quad T(3) = \boxed{1} \boxed{1} \boxed{2} \boxed{5}.$$

Note that the weight μ of $[T(j)]$ is a partition, while none of the weights of the individual tableaux $T(j)$ are partitions.

Given $[\lambda(j)] \in \mathcal{P}_n[r]$ and any composition ω of n , we define the *multipartition Kostka number* $K_{[\lambda(j)]\omega}$ to be the total number of semistandard Young r -multitableaux of shape $[\lambda(j)]$ and weight ω . The following relates the Kostka number for an r -multipartition to the Kostka number for the individual partitions.

Proposition 2.1. *For any composition v of length l and size n , and any $[\lambda(j)] \in \mathcal{P}_n[r]$, we have*

$$K_{[\lambda(j)]v} = \sum_{\substack{\sum_j \omega(j) = v \\ |\omega(j)| = |\lambda(j)|}} \prod_{j=1}^r K_{\lambda(j)\omega(j)},$$

where the sum is over all possible ways to choose r l -tuples $\omega(j)$, $1 \leq j \leq r$, of non-negative integers, where the sum of the $\omega(j)$ coordinate-wise is v , and the size of $\omega(j)$ taken as a composition is the same as the size of $\lambda(j)$.

Proof. We may count possible semistandard Young r -multitableaux of weight v and shape $[\lambda(j)]$ as follows. For each j , we choose the weight $\omega(j)$ of the tableau of shape $\lambda(j)$. We must choose these $\omega(j)$ so that the coordinate-wise sum is exactly

the weight v for the whole r -multitableau. Now $K_{\lambda(j)\omega(j)}$ is the total number of semistandard Young tableaux of shape $\lambda(j)$ and weight $\omega(j)$, so that once we fix the individual weights $\omega(j)$, the total number of r -multitableaux with this prescription is the product $\prod_{j=1}^r K_{\lambda(j)\omega(j)}$. The formula follows. \square

Next we observe that, like in the case for partitions, the Kostka number for a multipartition is invariant under permuting the parts of the weight. This is precisely why we may restrict our attention to the weight being a partition.

Corollary 2.2. *For any $[\lambda(j)] \in \mathcal{P}_n[r]$, and any composition v of size n , let μ be the unique partition obtained by permuting the parts of v . Then $K_{[\lambda(j)]v} = K_{[\lambda(j)]\mu}$.*

Proof. If v has length l , let $\sigma \in S_l$ be the permutation which, when applied to the parts of v , gives μ . If we use Proposition 2.1, and we apply σ to every l -tuple $\omega(j)$ in the sum when computing $K_{[\lambda(j)]v}$, then we obtain every l -tuple $\omega'(j)$ which would appear in the sum when computing $K_{[\lambda(j)]\mu}$. Since Kostka numbers for partitions are invariant under permutation of the parts of the weight, then we always have $K_{\lambda(j)\omega(j)} = K_{\lambda(j)\omega'(j)}$ when $\omega'(j)$ is the result of permuting the parts of $\omega(j)$ by σ . It follows that $K_{[\lambda(j)]v} = K_{[\lambda(j)]\mu}$. \square

The following result gives a precise condition for when the Kostka number for an r -multipartition is nonzero, and reduces to the case of partitions.

Lemma 2.3. *Suppose $[\lambda(j)] \in \mathcal{P}_n[r]$ and $\mu \in \mathcal{P}_n$. Then $K_{[\lambda(j)]\mu} > 0$ if and only if $\tilde{\lambda} \trianglerighteq \mu$.*

Proof. First suppose that $\tilde{\lambda}$ does not dominate μ (that is, suppose $\tilde{\lambda} \trianglerighteq \mu$ does not hold), but there does exist a semistandard Young r -multitableau $[T(j)]$ of shape $[\lambda(j)]$ and weight μ . Then for some $m \geq 1$, we have $\sum_{i \leq m} \tilde{\lambda}_i < \sum_{i \leq m} \mu_i$, which means

$$\sum_{j=1}^r \sum_{i \leq m} \lambda(j)_i < \sum_{i \leq m} \mu_i. \tag{2.5}$$

Since the entries in each column of each $T(j)$ strictly increase, then all entries 1 through m must appear in the first m rows of each $T(j)$. However, the inequality (2.5) says that there are more entries 1 through m than there are boxes in the first m rows of $[T(j)]$, giving a contradiction.

Now assume $\tilde{\lambda} \trianglerighteq \mu$. Then $K_{\tilde{\lambda}\mu} > 0$, and so there is a semistandard Young tableau T of shape $\tilde{\lambda}$ and weight μ . Given such a T , we may construct a semistandard Young r -multitableau $[T(j)]$ of shape $[\lambda(j)]$ and weight μ as follows. Each column of $\tilde{\lambda}$ is the same length of some column of some $\lambda(j)$, and we may take each column of T , and make it a column of some $T(j)$. If some column in T is to the left of another column in T , we need only make sure that if these columns are in the same $T(j)$, that the first is still to the left of the second. This guarantees that each $T(j)$ is a semistandard Young tableau, since the entries of each column strictly increase going down, as they did in T , and the entries in each row of $T(j)$ weakly increase to the right, since each column is arranged in an order so that weak increasing is preserved

from T . Thus any $[T(j)]$ created in this way is a semistandard Young r -multitableau of shape $[\lambda(j)]$ and weight μ , so that $K_{[\lambda(j)]\mu} > 0$. \square

The construction of the r -multitableau $[T(j)]$ from the tableau T in the end of the proof above will be used again, and so we give an example now.

Example 2.4. Let $\mu = (7, 6, 4, 3, 1, 1)$, and define $[\lambda(j)]$ by $\lambda(1) = (4, 4, 3)$, $\lambda(2) = (3, 3, 1)$, and $\lambda(3) = (3, 1)$. Then $\tilde{\lambda} = (10, 8, 4)$. One semistandard Young tableau T of shape $\tilde{\lambda}$ and weight μ is

$$T = \begin{array}{|c|c|c|c|c|c|c|c|c|c|} \hline 1 & 1 & 1 & 1 & 1 & 1 & 1 & 2 & 2 & 2 \\ \hline 2 & 2 & 2 & 3 & 3 & 3 & 3 & 3 & 4 & \\ \hline 4 & 4 & 5 & 6 & & & & & & \\ \hline \end{array}.$$

By choosing columns of T to go in different positions of an r -multitableau, while still preserving the order of entries, we can construct (at least) two different semistandard r -multitableau, $[T(j)]$ and $[T'(j)]$, of shape $[\lambda(j)]$ and weight μ , which we may define as

$$T(1) = \begin{array}{|c|c|c|c|} \hline 1 & 1 & 1 & 2 \\ \hline 2 & 2 & 3 & 4 \\ \hline 4 & 4 & 6 & \\ \hline \end{array}, \quad T(2) = \begin{array}{|c|c|c|} \hline 1 & 1 & 1 \\ \hline 2 & 3 & 3 \\ \hline 5 & & \\ \hline \end{array}, \quad T(3) = \begin{array}{|c|c|c|} \hline 1 & 2 & 2 \\ \hline 3 & & \\ \hline \end{array},$$

and

$$T'(1) = \begin{array}{|c|c|c|c|} \hline 1 & 1 & 1 & 1 \\ \hline 2 & 2 & 3 & 3 \\ \hline 4 & 5 & 6 & \\ \hline \end{array}, \quad T'(2) = \begin{array}{|c|c|c|} \hline 1 & 1 & 1 \\ \hline 2 & 3 & 3 \\ \hline 4 & & \\ \hline \end{array}, \quad T'(3) = \begin{array}{|c|c|c|} \hline 2 & 2 & 2 \\ \hline 4 & & \\ \hline \end{array}.$$

3 Multiplicity One for Partitions

Berenshteĭn and Zelevinskĭĭ [2] give precise conditions on partitions λ, μ , which are equivalent to the statement that $K_{\lambda\mu} = 1$. We note that there is a very slight typographical error in the subscripts of [2, Theorem 1.5] which we correct below.

Theorem 3.1 (Berenshteĭn and Zelevinskĭĭ). *Let $\lambda, \mu \in \mathcal{P}_n$, and suppose $\ell(\mu) = l$. Then $K_{\lambda\mu} = 1$ if and only if there exists a choice of indices $0 = i_0 < i_1 < \dots < i_t = l$ such that, for $k = 1, \dots, t$, the partitions*

$$\lambda^k = (\lambda_{i_{k-1}+1}, \lambda_{i_{k-1}+2}, \dots, \lambda_{i_k}) \quad \text{and} \quad \mu^k = (\mu_{i_{k-1}+1}, \mu_{i_{k-1}+2}, \dots, \mu_{i_k}),$$

where we define $\lambda_i = 0$ if $i > \ell(\lambda)$, satisfy the following:

- (1) $\lambda^k \supseteq \mu^k$ (so $|\lambda^k| = |\mu^k|$), and
- (2) $\lambda_{i_{k-1}+1} = \lambda_{i_{k-1}+2} = \dots = \lambda_{i_k-1}$, or $\lambda_{i_{k-1}+1} > \lambda_{i_{k-1}+2} = \lambda_{i_{k-1}+3} = \dots = \lambda_{i_k}$.

As our main result in Section 5 is a generalization of and heavily dependent on Theorem 3.1, we give a tableau-theoretic proof of Theorem 3.1 now. We begin with the following special cases.

Lemma 3.2. *Let $\lambda, \mu \in \mathcal{P}_n$ with $\ell(\mu) = l$, $\ell(\lambda) \leq l$, $\lambda \supseteq \mu$, and $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_l)$, with $\lambda_i = 0$ if $i > \ell(\lambda)$. Suppose that either:*

- (1) $\lambda_1 = \lambda_2 = \dots = \lambda_l,$
- (2) $\lambda_1 > \lambda_2 = \lambda_3 = \dots = \lambda_l,$ or
- (3) $\lambda_1 = \lambda_2 = \dots = \lambda_{l-1} > \lambda_l.$

Then $K_{\lambda\mu} = 1$. In case (1), or in case (2) when $\lambda_1 - \lambda_2 = 1$, or in case (3) when $\lambda_{l-1} - \lambda_l = 1$, we must also have $\mu = \lambda$.

Proof. Since $\lambda \geq \mu$, there is at least one semistandard Young tableau of shape λ and weight μ . Let T be such a tableau.

In case (1), we have $\ell(\lambda) = \ell(\mu)$, and the only case for T is that every column of T has entries 1 through l in sequential order, since entries must strictly increase from top to bottom in columns. Thus $K_{\lambda\mu} = 1$ and $\mu = \lambda$.

In case (2), if we have $\lambda_2 = 0$, then $\lambda = (\lambda_1)$, and we must have $K_{\lambda\mu} = 1$ since any semistandard Young tableau T of shape a single row must have entries in weakly increasing order. So assume $\lambda_2 > 0$, in which case $\ell(\lambda) = \ell(\mu)$. Since $\ell(\mu) = l$ is the number of rows of T , then the first λ_2 columns of T must have exactly the entries 1 through l in that order. The remaining $\lambda_1 - \lambda_2$ entries must be in weakly increasing order in row 1 of T . For example, if $\lambda = (6, 3, 3)$ and $\mu = (5, 4, 3)$, then the first 3 columns of the tableau must have entries 1, 2, 3, and the remaining entries must be in row 1 in a fixed order:

$$\begin{array}{|c|c|c|} \hline 1 & 1 & 1 \\ \hline 2 & 2 & 2 \\ \hline 3 & 3 & 3 \\ \hline \end{array} \mapsto \begin{array}{|c|c|c|c|c|c|} \hline 1 & 1 & 1 & 1 & 1 & 2 \\ \hline 2 & 2 & 2 & & & \\ \hline 3 & 3 & 3 & & & \\ \hline \end{array} .$$

There is thus only one such T , and $K_{\lambda\mu} = 1$. If $\lambda_1 - \lambda_2 = 1$, then the only entry that can appear in the right-most entry of the first row of T is 1, in order for the weight μ to be a partition. Thus $\mu = \lambda$ in this case.

In case (3), first consider the case that $\lambda_l = 0$, so that $\ell(\lambda) = l - 1$. In this case, each column of T is missing exactly one of the entries 1 through l . Since the entries of each column strictly increase from top to bottom, knowing the missing entry is the same as knowing the column entries. Since also row entries weakly increase from left to right, then given columns of length $l - 1$ with entries from 1 through l , the columns can only be arranged in one way to form T , which is in such a way that the missing entries from each column weakly decrease from left to right. In other words, T is uniquely determined from λ and μ , and $K_{\lambda\mu} = 1$. For example, if $\lambda = (5, 5, 5, 5)$ and $\mu = (5, 4, 4, 4, 3)$, then every column has a 1, one column each is missing a 2, 3, and 4, and one column is missing a 5. The only such T with this property is

$$\begin{array}{|c|c|c|c|c|} \hline 1 & 1 & 1 & 1 & 1 \\ \hline 2 & 2 & 2 & 2 & 3 \\ \hline 3 & 3 & 3 & 4 & 4 \\ \hline 4 & 4 & 5 & 5 & 5 \\ \hline \end{array} ,$$

where the missing entries from each column, from left to right, are 5, 5, 4, 3, and 2. Note that we cannot have $\lambda_{l-1} - \lambda_l = 1$ and $\ell(\mu) = l$ in this case. Now consider the

case that $\lambda_l > 0$. The first λ_l columns of λ are then of length l , and so these columns of T must have the entries 1 through l from top to bottom. If we remove these columns from λ and these entries from μ , we are left with a partition of length $l - 1$ with all equal parts, and now the case just covered (case (3) with $\lambda_l = 0$) applies. Thus $K_{\lambda\mu} = 1$. If $\lambda_{l-1} - \lambda_l = 1$, then again the entries of the first λ_l columns of T must be 1 through l from top to bottom, and the only choice for the last column of T to have the weight μ a partition, is to have the entries 1 through $l - 1$ from top to bottom. This implies $\mu = \lambda$. \square

We will also need the following result.

Lemma 3.3. *Let $\lambda, \mu \in \mathcal{P}_n$ such that $K_{\lambda\mu} = 1$ and $\ell(\lambda) = h \leq \ell(\mu) = l$. Then the unique semistandard Young tableau T of shape λ and weight μ has the following properties.*

- (1) *The first entry in each row i of T is i for $i = 1, \dots, h$.*
- (2) *If $h = l$, then every entry of row h of T is h . If $h < l$, then every entry greater than h in T is in row h of T , and further $\sum_{j=h+1}^l \mu_j < \lambda_h$.*

Proof. For (1), by way of contradiction suppose that i is minimal such that the first entry of row i of T is not i . If the first entry of row i is $j > i$, then this j entry does not have a $j - 1$ entry directly above it. Since $\mu_{j-1} \geq \mu_j$, then T must have a $j - 1$ entry with no j entry directly below it. In particular, such an entry has either no square below it, or has an entry greater than j below it. In a row of T with such an entry, consider the right-most such $j - 1$ entry. We may switch this $j - 1$ entry with the first j entry in row i , resulting in another distinct semistandard Young tableau of shape λ and weight μ , contradicting the fact that $K_{\lambda\mu} = 1$.

For statement (2), we consider the construction of T by greedily filling horizontal strips from bottom to top, as described in Section 2.2. Since $K_{\lambda\mu} = 1$, then this construction must yield the unique tableau T . By the construction, we begin by filling row h of T from right to left with the l entries, and moving to entries in shorter columns if we fill up row h , and then continuing with smaller entries on the next horizontal strip up. We note that by part (1), T must have the property that row h has first entry h , which means that when constructing T in this way, if $h = l$, then every entry of row h must be h , and if $h < l$ then we never fill up row h with entries greater than h . This implies that T has the property that all entries greater than h must be in row h , and these do not occupy all of row h . In other words, $\mu_{h+1} + \dots + \mu_l < \lambda_h$. \square

We remark that the proof of property (1) in Lemma 3.3 also implies that whenever $\lambda \triangleright \mu$, there always exists a semistandard Young tableau T of shape λ and weight μ such that each row i of T has first entry i , for $i = 1, \dots, h$, although we will not need this statement here.

Proof of Theorem 3.1. First we assume that $\lambda, \mu \in \mathcal{P}_n$ satisfy the conditions listed in Theorem 3.1. We have $\sum_{k=1}^t |\mu^k| = |\mu| = n$, $|\lambda^k| = |\mu^k|$ for $k = 1, \dots, t$, and $|\lambda| = n$.

It follows that each part of λ must be a part of some λ^k , otherwise $\sum_{k=1}^t |\lambda^k|$ is strictly less than n . In other words, the last index i_t must satisfy $i_t \geq \ell(\lambda)$, and thus $i_t = l = \ell(\mu) \geq \ell(\lambda)$. Also, since $\lambda^k \supseteq \mu^k$ for each k , it follows that $\lambda \supseteq \mu$, and so $K_{\lambda\mu} \geq 1$.

Consider a semistandard Young tableau T of shape λ and weight μ . Since entries strictly increase down columns of λ , any entry in the i th row of T must be at least i . Consider rows 1 through i_1 of T . Since any row past i_1 must have entries at least $i_1 + 1$, then all $|\mu^1|$ entries of value at most i_1 must occur in the first i_1 rows of T . Since $|\lambda^1| = |\mu^1|$, it follows that the *only* entries in the first i_1 rows of T can be 1 through i_1 . By induction, the only entries in rows $i_{k-1} + 1$ through i_k can be the values $i_{k-1} + 1$ through i_k . It follows that we may consider tableaux of shapes λ^k and weights μ^k independently, and that $K_{\lambda\mu} = \prod_{k=1}^t K_{\lambda^k\mu^k}$. We remark that this argument implies that this factorization result holds for *any* choice of indices, as long as $|\lambda^k| = |\mu^k|$ for each k . Now, it is enough to show that for $\lambda, \mu \in \mathcal{P}_n$, with $\ell(\mu) = l$, if $\lambda \supseteq \mu$ and either $\lambda_1 = \lambda_2 = \dots = \lambda_{l-1}$ or $\lambda_1 > \lambda_2 = \lambda_3 = \dots = \lambda_l$ (where $\lambda_i = 0$ if $i > \ell(\lambda)$), then $K_{\lambda\mu} = 1$. This is exactly what is proved in Lemma 3.2.

We now assume $K_{\lambda\mu} = 1$, and we prove there always exists a choice of indices $0 = i_0 < i_1 < i_2 < \dots < i_t = l$ with the desired properties. Throughout, we take T to be the unique semistandard Young tableau of shape λ and weight μ . We prove the statement by induction on n , where in the case $n = 1$, we have $\lambda = (1) = \mu$, and the result is immediate. So we assume $n > 1$, and that the statement holds for all positive integers less than n . We first consider the special cases that $\lambda = (1, 1, \dots, 1)$ or $\lambda = (n)$. In either case, note that if $K_{\lambda\mu} > 0$ then $K_{\lambda\mu} = 1$ since there is a single column or row, respectively. If $\lambda = (1, 1, \dots, 1)$, we must also have $\mu = (1, 1, \dots, 1)$ since entries must increase down the single column of T , and $\ell(\mu) = \ell(\lambda) = l$. Then take $t = 1$, so $\lambda^1 = \lambda$ and $\mu^1 = \mu$, and the desired conditions are satisfied. If $\lambda = (n)$, then μ may be any partition of n . Again take $t = 1$, so $\mu^1 = \mu$ and $\lambda^1 = (n, 0, \dots, 0)$, where the number of 0's is $l - 1$. The desired conditions are satisfied in this case as well, so we may assume that λ takes neither of these shapes. We now consider four scenarios in the induction, and we give an example in each to demonstrate the argument.

Let $\ell(\lambda) = h \leq \ell(\mu) = l$. We first consider the case that $\mu_{h+1} < \mu_h$ (which includes the case that $h = l$) and $\mu_h > 1$. For example, let $\lambda = (5, 3^3)$ and $\mu = (4^2, 3, 2, 1)$, so $h = 4 \leq l = 5$, and $\mu_h = 2 > \mu_{h+1} = 1$. Then we have

$$T = \begin{array}{|c|c|c|c|c|} \hline 1 & 1 & 1 & 1 & 2 \\ \hline 2 & 2 & 2 & & \\ \hline 3 & 3 & 3 & & \\ \hline 4 & 4 & 5 & & \\ \hline \end{array} .$$

Define new partitions λ^* and μ^* by $\lambda_i^* = \lambda_i - 1$ and $\mu_i^* = \mu_i - 1$ for $1 \leq i \leq h$, and $\mu_i^* = \mu_i$ for $i > h$. The assumption that $\mu_{h+1} < \mu_h$ guarantees that μ^* is indeed a partition, and we have $\lambda^*, \mu^* \in \mathcal{P}_{n-h}$. In our example, $\lambda^* = (4, 2^3)$ and $\mu^* = (3^2, 2, 1^2)$. Since $\lambda \supseteq \mu$, we also have $\lambda^* \supseteq \mu^*$. Consider any semistandard Young tableau T^* of shape λ^* and weight μ^* . Given T^* , add a box with an i entry at the beginning of row i for $i = 1, \dots, h$ (noting that some of these rows of T^* might be

initially length 0). Since the least possible entry in row i of a semistandard Young tableau is i , then this construction yields a semistandard Young tableau of shape λ and weight μ , which then must be T since $K_{\lambda\mu} = 1$, and note that T has the property that the first entry of each row i is i by Lemma 3.3. Consequently, T^* must be the unique semistandard Young tableau of shape λ^* and weight μ^* , since a distinct such tableau would yield a distinct tableau of shape λ and weight μ . Thus, $K_{\lambda^*\mu^*} = 1$, and we may apply the induction hypothesis to λ^* and μ^* . In the example, we have

$$T^* = \begin{array}{|c|c|c|c|} \hline 1 & 1 & 1 & 2 \\ \hline 2 & 2 & & \\ \hline 3 & 3 & & \\ \hline 4 & 5 & & \\ \hline \end{array} .$$

Let $0 = i_0^* < i_1^* < \dots < i_t^* = \ell(\mu^*)$ be the desired indices for λ^* and μ^* , which exist by the induction hypothesis, with the accompanying subpartitions λ^{*k} and μ^{*k} . In the example, we can take $i_1^* = 2$ and $i_2^* = 5$, with $\lambda^{*1} = (4, 2)$, $\mu^{*1} = (3, 3)$, $\lambda^{*2} = (2, 2, 0)$, and $\mu^{*2} = (2, 1, 1)$. Note that $\ell(\mu^*) = \ell(\mu)$ since $\mu_h > 1$. Now let $i_k = i_k^*$ for $k = 0, 1, \dots, t$, and consider the corresponding subpartitions λ^k and μ^k . In the example, this gives $\lambda^1 = (5, 3)$, $\mu^1 = (4, 4)$, $\lambda^2 = (3, 3, 0)$, and $\mu^2 = (2, 1, 1)$. Since $\lambda^{*k} \supseteq \mu^{*k}$, then it follows that $\lambda^k \supseteq \mu^k$ for $k = 1, \dots, t$. We also have either $\lambda_{i_{k-1}^*+1}^* = \lambda_{i_{k-1}^*+2}^* = \dots = \lambda_{i_k^*-1}^*$ or $\lambda_{i_{k-1}^*+1}^* > \lambda_{i_{k-1}^*+2}^* = \lambda_{i_{k-1}^*+3}^* = \dots = \lambda_{i_k^*}^*$ for $k = 1, \dots, t$. By Lemma 3.3(2), every entry greater than h is in row h of T . In the case that $k < t$, we must then have $i_k \leq h-1$, and so either $\lambda_{i_{k-1}^*+1} = \lambda_{i_{k-1}^*+2} = \dots = \lambda_{i_k-1}$ or $\lambda_{i_{k-1}^*+1} > \lambda_{i_{k-1}^*+2} = \lambda_{i_{k-1}^*+3} = \dots = \lambda_{i_k}$ by how we defined λ^* . The $k = t$ case follows from the assumption that $\mu_h > 1$, because if $\lambda_h = 1$ then $\ell(\lambda) = \ell(\mu)$ (since the last row of T would have no room for entries greater than h), and if $\lambda_h > 1$ then $\ell(\lambda) = \ell(\lambda^*)$. These statements imply the desired subpartition properties are preserved.

If $\mu_{h+1} < \mu_h$ and $\mu_h = 1$, then $\mu_{h+1} = 0$ and $h = l$, and $\lambda_h = 1$, which follows from Lemma 3.3(2) since all entries in row h of T must be h . As an example, take $\lambda = (4, 2, 1)$ and $\mu = (3^2, 1)$, which means we have

$$T = \begin{array}{|c|c|c|c|} \hline 1 & 1 & 1 & 2 \\ \hline 2 & 2 & & \\ \hline 3 & & & \\ \hline \end{array} .$$

Now let $\lambda^* = (\lambda_1, \dots, \lambda_{l-1})$ and $\mu^* = (\mu_1, \dots, \mu_{l-1})$, so in the example $\lambda^* = (4, 2)$ and $\mu^* = (3^2)$. Given any semistandard Young tableau of shape λ^* and weight μ^* , we may add a single box with entry l at the bottom to obtain a semistandard Young tableau of shape λ and weight μ . Since $K_{\lambda\mu} = 1$, the resulting tableau must be T . Thus we must have $K_{\lambda^*\mu^*} = 1$, since otherwise we could obtain more than one semistandard Young tableau of shape λ and weight μ . In the example,

$$T^* = \begin{array}{|c|c|c|c|} \hline 1 & 1 & 1 & 2 \\ \hline 2 & 2 & & \\ \hline \end{array} .$$

We take the indices obtained by applying the induction hypothesis to $\lambda^*, \mu^* \in \mathcal{P}_{n-1}$, and we add to it the index $i_t = l$ to obtain indices for λ and μ . In the example,

$i_1^* = 2$, with $\lambda^{*1} = \lambda^* = (4, 2)$ and $\mu^{*1} = \mu^* = (3, 3)$, and then $i_1 = 2, i_2 = 3$, with $\lambda^1 = (4, 2), \mu^1 = (3, 3)$, and $\lambda^2 = (1) = \mu^2$. Since we will always have $\lambda^t = (1) = \mu^t$, and $\lambda^{*k} = \lambda^k, \mu^{*k} = \mu^k$ for $k < t$, it follows that the resulting subpartitions have the desired properties.

Now consider the case that $\mu_{h+1} = \mu_h$ (so $\ell(\lambda) = h < \ell(\mu) = l$) and either $l > h + 1$ or $\lambda_h = \lambda_{h-1}$. Take as an example $\lambda = (5, 3^3)$ and $\mu = (4^2, 3, 1^3)$, so $h = 4$ and $l = 6$, and $\lambda_4 = \lambda_3 = 1$. In this case,

$$T = \begin{array}{|c|c|c|c|c|} \hline 1 & 1 & 1 & 1 & 2 \\ \hline 2 & 2 & 2 & & \\ \hline 3 & 3 & 3 & & \\ \hline 4 & 5 & 6 & & \\ \hline \end{array} .$$

By Lemma 3.3(2), all entries greater than h must be in row h of T . Now define λ^* and μ^* by $\lambda_i^* = \lambda_i$ if $i < h$, and $\lambda_h^* = \lambda_h - \mu_l$, and $\mu_i^* = \mu_i$ if $i < l$, and $\mu_i^* = 0$ if $i = l$. Then $\lambda^* = (5, 3^2, 2)$ and $\mu^* = (4^2, 3, 1^2)$ in the example. Note that $\lambda_h^* > 0$ by Lemma 3.3(1), since row h of T has as its first entry $h < l$. Now $\lambda^*, \mu^* \in \mathcal{P}_{n-\mu_l}$, and $\lambda^* \supseteq \mu^*$ since $\lambda \supseteq \mu$. If T^* is any semistandard Young tableau of shape λ^* and weight μ^* , then we can add μ_l boxes with entries of l to row h of T^* , and we obtain a semistandard Young tableau of shape λ and weight μ . This must be T , since $K_{\lambda\mu} = 1$, and then we must also have $K_{\lambda^*\mu^*} = 1$. In our example,

$$T^* = \begin{array}{|c|c|c|c|c|} \hline 1 & 1 & 1 & 1 & 2 \\ \hline 2 & 2 & 2 & & \\ \hline 3 & 3 & 3 & & \\ \hline 4 & 5 & & & \\ \hline \end{array} .$$

Let $0 = i_0^* < i_1^* < \dots < i_t^* = \ell(\mu^*) = l - 1$ be the indices which exist by the induction hypothesis, with the subpartitions λ^{*k} and μ^{*k} with the desired properties. We may take $i_1^* = 3$ and $i_2^* = 5$, with $\lambda^{*1} = (5, 3, 3), \mu^{*1} = (4, 4, 3), \lambda^{*2} = (2, 0)$, and $\mu^{*2} = (1, 1)$ in the example. Define $i_k = i_k^*$ if $k < t$, and $i_t = i_t^* + 1 = l$. In the example, this gives $i_1 = 3, i_2 = 6, \lambda^1 = (5, 3, 3) = \lambda^{*1}, \mu^1 = (4, 4, 3) = \mu^{*1}, \lambda^2 = (3, 0, 0)$, and $\mu^2 = (1, 1, 1)$. In general we have $\lambda^k = \lambda^{*k}$ and $\mu^k = \mu^{*k}$ if $k < t$, so the desired properties hold. When $k = t$, then $\lambda^k \supseteq \mu^k$, and the fact that λ^k is of the desired shape follows from the assumption that either $l > h + 1$, in which case $i_{t-1} = h - 1$ necessarily, or $\lambda_h = \lambda_{h-1}$ with $l = h + 1$, in which case $i_t^* = h$ and $i_t = h + 1 = l$.

Finally, consider the case with $\mu_{h+1} = \mu_h, l = h + 1$, and $\lambda_h < \lambda_{h-1}$. For example, let $\lambda = (5, 3^2, 2)$ and $\mu = (4^2, 3, 1^2)$, so $h = 4, l = 5$, and

$$T = \begin{array}{|c|c|c|c|c|} \hline 1 & 1 & 1 & 1 & 2 \\ \hline 2 & 2 & 2 & & \\ \hline 3 & 3 & 3 & & \\ \hline 4 & 5 & & & \\ \hline \end{array} .$$

Assume that in row $h - 1$ of T , there are some h entries, and we will obtain a contradiction. This implies that the right-most entry in row $h - 1$ is h , since all μ_l of

the $l = h + 1$ entries in T are in row h of T , by Lemma 3.3. Consider the left-most l entry in row h of T . There cannot be an $l - 1 = h$ entry directly above this entry, since then all entries to the right of it would also be h , which would imply $\mu_h > \mu_{h+1}$, a contradiction. Then we can exchange the left-most $l = h + 1$ entry in row h with the right-most h entry in row $h - 1$ of T to obtain a distinct semistandard Young tableau T' of shape λ and weight μ , contradicting $K_{\lambda\mu} = 1$. Thus, there are no h entries in row $h - 1$ of T , and thus no h entries above row h of T by the greedy filling of horizontal strips, since $\lambda_h < \lambda_{h-1}$, and if there are other h entries, there must be some in row $h - 1$. In particular, we have $\lambda_h = \mu_h + \mu_{h+1}$. Now define λ^* by $\lambda_i^* = \lambda_i$ if $i < h$ and $\lambda_h^* = 0$, and we define μ^* by $\mu_i^* = \mu_i$ if $i < h$ and $\mu_h^* = \mu_{h+1}^* = 0$. In the example, $\lambda^* = (5, 3^2)$ and $\mu^* = (4^2, 3)$. In any semistandard Young tableau T^* of shape λ^* and weight μ^* , we can always add a row of length λ_h containing μ_h entries of h and μ_{h+1} entries of $h + 1$, since T^* contains entries $h - 1$ or lower. As in the previous cases, the resulting tableau must be T since $K_{\lambda\mu} = 1$, and so T^* is the unique semistandard Young tableau of shape λ^* and weight μ^* , and $K_{\lambda^*\mu^*} = 1$. In our example,

$$T^* = \begin{array}{|c|c|c|c|c|} \hline 1 & 1 & 1 & 1 & 2 \\ \hline 2 & 2 & 2 & & \\ \hline 3 & 3 & 3 & & \\ \hline \end{array} .$$

We may then apply the induction hypothesis to λ^* and μ^* . Take the indices for λ^* and μ^* from the induction hypothesis, and define these to be i_0 through $i_{t-1} = h - 1$ for λ and μ , and $i_t = h + 1 = l$. So $\lambda^{*k} = \lambda^k$ and $\mu^{*k} = \mu^k$ for $k \leq t - 1$, and $\lambda^t = (\lambda_h, 0)$, $\mu^t = (\mu_h, \mu_{h+1})$. In the example, we may take $i_1^* = 3$, so $i_1 = 2$ and $i_2 = 5$, with $\lambda^{*1} = \lambda^1 = (5, 3, 3)$, $\mu^{*1} = \mu^1 = (4, 4, 3)$, and $\lambda^2 = (2, 0)$, $\mu^2 = (1, 1)$. By the induction hypothesis, and since $\lambda^t = (\lambda_h, 0)$ always satisfies $\lambda^t \succeq \mu^t = (\mu_h, \mu_{h+1})$, this choice gives the desired indices for λ and μ . This completes the induction. \square

It follows from Theorem 3.1, or from direct observation, that for any partition λ we have $K_{\lambda\lambda} = 1$. The next result gives exactly which partitions λ satisfy $K_{\lambda\mu} = 1$ for only $\mu = \lambda$.

Corollary 3.4 (Gates, Goldman, and Vinroot). *Let $\lambda \in \mathcal{P}$. The only $\mu \in \mathcal{P}$ which satisfies $K_{\lambda\mu} = 1$ is $\mu = \lambda$ if and only if $\lambda_i - \lambda_{i+1} \leq 1$ for all i (where $\lambda_i = 0$ if $i > \ell(\lambda)$).*

The proof of Corollary 3.4 as it appears in [7, Corollary 5.1] follows from an enumeration of those μ which satisfy $K_{\lambda\mu} = 1$. We could give another proof of Corollary 3.4 here as a corollary of Theorem 3.1 and Lemma 3.2. Instead, we generalize the statement in Corollary 5.4, with a proof independent of Corollary 3.4.

Our main goal is to give conditions on an r -multipartition $[\lambda(j)]$ and a partition μ which are equivalent to $K_{[\lambda(j)]\mu} = 1$, and to generalize Corollary 3.4 to multipartitions. Before accomplishing this goal, we make a departure into representation theory to find motivation to study these Kostka numbers.

4 Permutation Characters of Wreath Products

Let S_n denote the symmetric group on n elements. For any group G , and any subgroup of permutations $P \leq S_n$, we let $G \wr P$ denote the semidirect product $G^n \rtimes P$, where G^n is the direct product of n copies of G , and P acts by inverse permutation of indices on elements in G^n . That is, if $g = (g_1, \dots, g_n) \in G^n$ and $\tau \in P$, then the action of τ on g is given by

$$\tau g = (g_{\tau^{-1}(1)}, \dots, g_{\tau^{-1}(n)}).$$

If $P = S_n$, then $G \wr S_n$ is the standard wreath product.

Let A and B be finite groups, with complex representations (π, V) and (ρ, W) , with characters χ and η , respectively. We let $\pi \odot \rho$ denote the external tensor product of representations, which is a representation of $A \times B$ acting on the space $V \otimes W$, with character given by $(\chi \odot \eta)(a, b) = \chi(a)\eta(b)$. In particular, $\bigodot_{i=1}^n \pi = \pi^{\odot n}$ is a representation of A^n acting on $\bigotimes_{i=1}^n V = V^{\otimes n}$, with character $\bigodot_{i=1}^n \chi = \chi^{\odot n}$. If (π, V) and (ρ, W) are both representations of A , then we let $\pi \otimes \rho$ denote the internal tensor product of representations, which is a representation of A acting on $V \otimes W$, with character given by $(\chi \otimes \eta)(a) = \chi(a)\eta(a)$.

We let $\mathbf{1}$ denote the trivial character of a finite group. For any finite group A , let $\text{Irr}(A)$ denote the collection of irreducible complex characters of A . Given any character χ , we let $\chi(1)$ denote its degree. If χ and η are complex-valued class functions of A , let $\langle \chi, \eta \rangle = 1/|A| \sum_{a \in A} \chi(a)\eta(a)$ be the standard inner product. For a subgroup $D \leq A$, and a character ξ of D , we let $\text{Ind}_D^A(\xi)$ denote the induced character from D to A .

As in Section 2.2, for any $\lambda \in \mathcal{P}_n$, we let π^λ be the irreducible representation of S_n associated with λ , with character χ^λ , in such a way that $\chi^{(n)} = \mathbf{1}$ and $\chi^{(1,1,\dots,1)}$ is the sign character.

4.1 Irreducible characters of wreath products

Let G be any finite group, and let $G_n = G \wr S_n$. The irreducible complex characters of G_n were first described by Specht [22], and the description we give in this section and the next follows [12, Section 4.3] and [16, Chapter I, Appendix B], and we have also found [17, Section 4] helpful. Another approach to the representation theory of wreath products is given in [4].

If (ϱ, V) is a representation of G , then $\varrho^{\odot n}$ is a representation of G^n acting on $V^{\otimes n}$. We extend $\varrho^{\odot n}$ to a representation $\underline{\varrho}^{\odot n}$ of $G_n = G \wr S_n$ by defining, for $g = (g_1, \dots, g_n) \in G^n$, $\tau \in S_n$, and $v_i \in V$,

$$(\underline{\varrho}^{\odot n})(g, \tau)(v_1 \otimes \dots \otimes v_n) = (\varrho(g_1)v_{\tau^{-1}(1)} \otimes \dots \otimes \varrho(g_n)v_{\tau^{-1}(n)}).$$

Given the representation π^λ , $\lambda \in \mathcal{P}_n$, of S_n , extend π^λ to a representation $\underline{\pi}^\lambda$ of G_n trivially, that is,

$$\underline{\pi}^\lambda(g, \tau) = \pi^\lambda(\tau), \quad \text{for } g \in G^n, \tau \in S_n.$$

Then, given a representation ϱ of G and $\lambda \in \mathcal{P}_n$, define the representation $\varrho \wr \lambda$ of G_n by

$$\varrho \wr \lambda = (\varrho^{\odot n}) \otimes \underline{\pi}^\lambda,$$

and if ψ is the character of ϱ , we let $\psi \wr \lambda$ denote the character of $\varrho \wr \lambda$.

Suppose that G has $r = |\text{Irr}(G)|$ irreducible characters, and consider the set $\mathcal{P}_n[r]$ of r -multipartitions of n . We would like to associate a partition to each $\psi \in \text{Irr}(G)$, and while we could label each irreducible character of G as ψ_j for $1 \leq j \leq r$ and associate $\lambda(j)$ with ψ_j , we instead write $\lambda(\psi)$ for this partition in the general case for the sake of clarity. That is, for the parameterization which we will describe, we write $[\lambda(\psi)] \in \mathcal{P}_n[r]$ with the understanding that each $\psi \in \text{Irr}(G)$ is labeled by $1 \leq j \leq |\text{Irr}(G)|$ in some order, and that $\lambda(\psi)$ is taken as $\lambda(j)$, with ψ labeled by j .

Given any $[\lambda(\psi)] \in \mathcal{P}_n[|\text{Irr}(G)|]$, define the subgroup $G[\lambda(\psi)]$ of G_n by

$$G[\lambda(\psi)] = \prod_{\psi \in \text{Irr}(G)} G_{|\lambda(\psi)|} = \prod_{\psi \in \text{Irr}(G)} (G \wr S_{|\lambda(\psi)|}).$$

Then we have that

$$\bigodot_{\psi \in \text{Irr}(G)} \psi \wr \lambda(\psi) \quad \text{is a character of } G[\lambda(\psi)].$$

We may now give the parameterization of irreducible characters of G_n .

Theorem 4.1 (Specht [22]). *The irreducible complex characters of G_n may be parameterized by $\mathcal{P}_n[|\text{Irr}(G)|]$, where for each $[\lambda(\psi)] \in \mathcal{P}_n[|\text{Irr}(G)|]$, the irreducible character $\eta^{[\lambda(\psi)]}$ is given by*

$$\eta^{[\lambda(\psi)]} = \text{Ind}_{G[\lambda(\psi)]}^{G_n} \left(\bigodot_{\psi \in \text{Irr}(G)} \psi \wr \lambda(\psi) \right).$$

The degree of $\eta^{[\lambda(\psi)]}$ is given by

$$\eta^{[\lambda(\psi)]}(1) = n! \prod_{\psi \in \text{Irr}(H)} \frac{(\psi(1)^{|\lambda(\psi)|}) (\chi^{\lambda(\psi)}(1))}{|\lambda(\psi)|!}.$$

4.2 Symmetric functions

Macdonald [16, Chapter 1, Appendix B] gives a method of describing the irreducible characters of G_n using symmetric function theory, which is an extension of the description of the character theory of the symmetric groups using symmetric functions [16, Chapter 1.8]. Another detailed treatment of symmetric functions associated with wreath products is given by Ingram, Jing, and Stitzinger [11].

For each $\psi \in \text{Irr}(G)$, let $Y_\psi = \{y_{i\psi} \mid i \geq 1\}$ be an infinite set of indeterminates, where any two $y_{i\psi}$ commute. For any partition λ , let $s_\lambda(Y_\psi)$ be the Schur symmetric function in these variables. For any sequence $[\lambda(\psi)] \in \mathcal{P}_n[|\text{Irr}(G)|]$, define

$$s_{[\lambda(\psi)]} = \prod_{\psi \in \text{Irr}(G)} s_{\lambda(\psi)}(Y_\psi).$$

Let $\Lambda_n = \mathbb{C}\text{-span}\{s_{[\lambda(\psi)]} \mid [\lambda(\psi)] \in \mathcal{P}_n[[\text{Irr}(G)]]\}$ and $\Lambda = \bigoplus_n \Lambda_n$. Then Λ is a graded \mathbb{C} -algebra, where multiplication is standard multiplication of symmetric functions coming from polynomial multiplication. Let $R(G_n)$ be defined as the space of \mathbb{C} -valued class functions of G_n , and let $R = \bigoplus_n R(G_n)$. For any $m, n \geq 1$, we may embed $G_m \times G_n$ in G_{m+n} by embedding $S_m \times S_n$ in S_{m+n} as we did in Section 2.2, and we identify $G_m \times G_n$ as a subgroup of G_{m+n} in this way. For $\alpha \in R(G_n)$ and $\beta \in R(G_m)$, define the product $\alpha\beta \in R(G_{m+n})$ by the induced class function

$$\alpha\beta = \text{Ind}_{G_m \times G_n}^{G_{m+n}}(\alpha \times \beta).$$

This multiplication makes R a graded \mathbb{C} -algebra. We only need parts of the main result in [16, Chapter 1, Appendix B], and the following is a portion of [16, I.B, (9.7)].

Theorem 4.2. *Let $\eta^{[\lambda(\psi)]}$ be the irreducible character of $G_n = G \wr S_n$ corresponding to $[\lambda(\psi)] \in \mathcal{P}_n[[\text{Irr}(G)]]$. There is an isomorphism of graded \mathbb{C} -algebras*

$$\text{ch} : R \rightarrow \Lambda,$$

which we may define by $\text{ch}(\eta^{[\lambda(\psi)]}) = s_{[\lambda(\psi)]}$, and extend linearly. In particular, for any $\alpha, \beta \in R$, $\text{ch}(\alpha\beta) = \text{ch}(\alpha)\text{ch}(\beta)$.

For any $|\text{Irr}(G)|$ -multipartition $[\lambda(\psi)]$, define the *height* of the multipartition as

$$\text{ht}([\lambda(\psi)]) = \max\{\ell(\lambda(\psi)) \mid \psi \in \text{Irr}(G)\}.$$

In particular, $\text{ht}([\gamma(\psi)]) = 1$ if and only if each $\gamma(\psi)$ is either the empty partition or has a single part (and at least one $\gamma(\psi)$ is not the empty partition).

For any $m \geq 1$ and any subset $\mathcal{S} \subseteq \text{Irr}(G)$, define the element $H_m^{(\mathcal{S})} \in \Lambda_m$ as

$$H_m^{(\mathcal{S})} = \sum_{\substack{|\gamma(\psi)|=m \\ \text{ht}([\gamma(\psi)])=1 \\ \gamma(\psi)=(0) \text{ if } \psi \notin \mathcal{S}}} s_{[\gamma(\psi)]}.$$

For any partition $\mu = (\mu_1, \mu_2, \dots, \mu_l)$ of n , define $H_\mu^{(\mathcal{S})} \in \Lambda_n$ as

$$H_\mu^{(\mathcal{S})} = H_{\mu_1}^{(\mathcal{S})} H_{\mu_2}^{(\mathcal{S})} \dots H_{\mu_l}^{(\mathcal{S})}.$$

We now prove a generalization of (2.1) in the algebra Λ . This result is implied by the wreath product versions of the Littlewood-Richardson rule proved by Ingram, Jing, and Stitzinger (via symmetric functions) [11, Theorem 4.7], and by Stein [23, Theorem 4.5] (proved and generalized in a categorical setting). The result is equivalent to [11, Corollary 4.6], but we give a proof of the result here for the purposes of self-containment.

Lemma 4.3. *For any partition μ of n , we have*

$$H_\mu^{(\mathcal{S})} = \sum_{\substack{|\lambda(\psi)|=n \\ \lambda(\psi)=(0) \text{ if } \psi \notin \mathcal{S}}} K_{[\lambda(\psi)]\mu} s_{[\lambda(\psi)]}.$$

Proof. Each factor $H_{\mu_i}^{(\mathcal{S})}$ consists of summands $s_{[\gamma(\psi)]}$, such that each partition $\gamma(\psi)$ is either empty or a single row, so write $\gamma(\psi) = (\gamma(\psi)_1)$, where we take $\gamma(\psi)_1 = 0$ when $\gamma(\psi) = (0)$. Then we have

$$s_{[\gamma(\psi)]} = \prod_{\psi \in \text{Irr}(G)} s_{(\gamma(\psi)_1)}(Y_\psi) = \prod_{\psi \in \text{Irr}(G)} h_{\gamma(\psi)_1}(Y_\psi),$$

where h_k is the complete symmetric function. Now we have

$$\begin{aligned} H_\mu^{(\mathcal{S})} &= \prod_{i=1}^l H_{\mu_i}^{(\mathcal{S})} = \prod_{i=1}^l \left(\sum_{\substack{|\gamma^{(i)}(\psi)| = \mu_i \\ \text{ht}([\gamma^{(i)}(\psi)]) = 1 \\ \gamma^{(i)}(\psi) = (0) \text{ if } \psi \notin \mathcal{S}}} \prod_{\psi \in \text{Irr}(G)} h_{\gamma^{(i)}(\psi)_1}(Y_\psi) \right) \\ &= \sum_{\substack{1 \leq i \leq l \\ |\gamma^{(i)}(\psi)| = \mu_i \\ \text{ht}([\gamma^{(i)}(\psi)]) = 1 \\ \gamma^{(i)}(\psi) = (0) \text{ if } \psi \notin \mathcal{S}}} \prod_{\psi \in \text{Irr}(G)} h_{\gamma^{(1)}(\psi)_1}(Y_\psi) \cdots h_{\gamma^{(l)}(\psi)_1}(Y_\psi) \\ &= \sum_{\substack{1 \leq i \leq l \\ |\gamma^{(i)}(\psi)| = \mu_i \\ \text{ht}([\gamma^{(i)}(\psi)]) = 1 \\ \gamma^{(i)}(\psi) = (0) \text{ if } \psi \notin \mathcal{S}}} \prod_{\psi \in \text{Irr}(G)} \sum_{\lambda(\psi) \in \mathcal{P}_{|\omega(\psi)|}} K_{\lambda(\psi)\omega(\psi)} s_{\lambda(\psi)}(Y_\psi), \end{aligned}$$

where we take $\omega(\psi)$ to be the composition $\omega(\psi) = (\gamma^{(1)}(\psi)_1, \dots, \gamma^{(l)}(\psi)_1)$, and we have applied (2.1). Note that we have $\sum_\psi \omega(\psi) = \mu \vdash n$. We may now rearrange sums and products to obtain

$$\begin{aligned} H_\mu^{(\mathcal{S})} &= \sum_{\substack{|\lambda(\psi)| = n \\ \lambda(\psi) = (0) \text{ if } \psi \notin \mathcal{S}}} \left(\sum_{\substack{\sum_\psi \omega(\psi) = \mu \\ |\omega(\psi)| = |\lambda(\psi)|}} \prod_{\psi \in \text{Irr}(G)} K_{\lambda(\psi)\omega(\psi)} \right) s_{[\lambda(\psi)]} \\ &= \sum_{\substack{|\lambda(\psi)| = n \\ \lambda(\psi) = (0) \text{ if } \psi \notin \mathcal{S}}} K_{[\lambda(\psi)]\mu} s_{[\lambda(\psi)]}, \end{aligned}$$

where we have applied Proposition 2.1 to obtain the last equality. □

4.3 Permutation characters of G_n on certain subgroups

If G is any finite group with $L \leq G$ a subgroup, then $L_n = L \wr S_n$ is a subgroup of G_n . Consider the case when $L \trianglelefteq G$ is normal with G/L abelian. If $\psi \in \text{Irr}(G)$ such that $L \subseteq \ker(\psi)$, then ψ can be viewed as a character of G/L , and in particular $\psi(1) = 1$. Suppose that $\eta^{[\lambda(\psi)]} \in \text{Irr}(G_n)$ such that $\lambda(\psi) = (0)$ if $L \not\subseteq \ker(\psi)$. Then it follows from Theorem 4.1 that

$$\eta^{[\lambda(\psi)]}(1) = n! \prod_{\psi \in \text{Irr}(G)} \frac{\chi^{\lambda(\psi)}(1)}{|\lambda(\psi)|!}.$$

The decomposition of the permutation character of G_n on L_n in this situation is as follows.

Proposition 4.4. *Let G be a finite group and $L \trianglelefteq G$ a normal subgroup such that G/L is abelian. Then*

$$\text{Ind}_{L_n}^{G_n}(\mathbf{1}) = \sum_{\substack{|\gamma(\psi)|=n \\ \text{ht}(\gamma(\psi))=1 \\ \gamma(\psi)=(0) \text{ if } L \not\subseteq \ker(\psi)}} \eta^{[\gamma(\psi)]}.$$

Proof. We first show that the character degrees on both sides are the same. The degree of $\text{Ind}_{L_n}^{G_n}(\mathbf{1})$ is $[G_n : L_n] = |G/L|^n$. Consider one character $\eta^{|\gamma(\psi)|}$ in the sum on the right. Since each $\gamma(\psi)$ has a single part or is empty, then $\chi^{\gamma(\psi)}(1) = 1$. Since $L \subseteq \ker(\psi)$ whenever $\gamma(\psi) \neq (0)$, and since we may think of ψ as a character of G/L , then the multipartitions we are considering may be thought of as $|\text{Irr}(G/L)|$ -multipartitions. We now have

$$\eta^{|\gamma(\psi)|}(1) = \frac{n!}{\prod_{\psi \in \text{Irr}(G/L)} |\gamma(\psi)|!},$$

which is a multinomial coefficient. In summing the degrees of these characters, we are summing over all possible multinomial coefficients of degree n with $|G/L|$ terms, which has sum $|G/L|^n$.

Since the degrees are equal, it is now enough to show that any character $\eta^{[\gamma(\psi)]}$ in the sum appears with nonzero multiplicity in the permutation character. Indeed, we will show that

$$\langle \text{Ind}_{L_n}^{G_n}(\mathbf{1}), \eta^{[\gamma(\psi)]} \rangle = 1. \tag{4.1}$$

From Theorem 4.1, we have that

$$\eta^{[\gamma(\psi)]} = \text{Ind}_{G[\gamma(\psi)]}^{G_n}(\xi^{[\gamma(\psi)]}) \quad \text{where} \quad \xi^{[\gamma(\psi)]} = \bigodot_{\psi \in \text{Irr}(G)} \psi \wr \gamma(\psi).$$

By Frobenius reciprocity, we have

$$\langle \text{Ind}_{L_n}^{G_n}(\mathbf{1}), \eta^{[\gamma(\psi)]} \rangle = \langle \text{Res}_{G[\gamma(\psi)]}(\text{Ind}_{L_n}^{G_n}(\mathbf{1})), \xi^{[\gamma(\psi)]} \rangle,$$

where Res is restriction of characters. By Mackey’s theorem, we have

$$\text{Res}_{G[\gamma(\psi)]}(\text{Ind}_{L_n}^{G_n}(\mathbf{1})) = \sum_{g \in [G[\gamma(\psi)] \backslash G_n / L_n]} \text{Ind}_{G[\gamma(\psi)] \cap gL_n g^{-1}}^{G[\gamma(\psi)]}(\mathbf{1}),$$

where the sum is over a set of double coset representatives. A brief consideration reveals that the only double coset representative is $g = 1 \in G_n$. Define

$$L[\gamma(\psi)] = G[\gamma(\psi)] \cap L_n = \prod_{\psi \in \text{Irr}(G)} (L \wr S_{|\gamma(\psi)|}).$$

By Mackey’s theorem and Frobenius reciprocity, we then have

$$\begin{aligned} \langle \text{Ind}_{L_n}^{G_n}(\mathbf{1}), \eta^{[\gamma(\psi)]} \rangle &= \langle \text{Ind}_{L[\gamma(\psi)]}^{G[\gamma(\psi)]}(\mathbf{1}), \xi^{[\gamma(\psi)]} \rangle \\ &= \langle \mathbf{1}, \text{Res}_{L[\gamma(\psi)]}(\xi^{[\gamma(\psi)]}) \rangle \\ &= \frac{1}{|L[\gamma(\psi)]|} \sum_{x \in L[\gamma(\psi)]} \xi^{[\gamma(\psi)]}(x). \end{aligned}$$

We have $\gamma(\psi) = (0)$ unless $L \subseteq \ker(\psi)$, and we have $\chi^{\gamma(\psi)}$ is trivial for every $\psi \in \text{Irr}(G)$. It follows directly from the definition that for every $\psi \in \text{Irr}(G)$ and every $(a, \tau) \in L_{|\gamma(\psi)|} = L \wr S_{|\gamma(\psi)|}$ that

$$(\psi \wr \gamma(\psi))(a, \tau) = 1.$$

Thus, for any $x \in L[\gamma(\psi)]$, we have $\xi^{[\gamma(\psi)]}(x) = 1$, and (4.1) follows. □

Note that if G/L is not abelian, then Proposition 4.4 does not hold, in that the decomposition cannot be multiplicity-free since the degrees will not match.

We now see multipartition Kostka numbers as multiplicities, in the following generalization of (2.3) to wreath products.

Theorem 4.5. *Let G be a finite group, $L \trianglelefteq G$ a normal subgroup such that G/L is abelian, $n \geq 1$, and $\mu = (\mu_1, \dots, \mu_l) \in \mathcal{P}_n$. Define $L_\mu = L \wr S_\mu = \prod_{i=1}^l (L \wr S_{\mu_i})$. Then the permutation character of G_n on L_μ decomposes as*

$$\text{Ind}_{L_\mu}^{G_n}(\mathbf{1}) = \sum_{\substack{|\lambda(\psi)|=n \\ \lambda(\psi)=(0) \text{ if } L \not\subseteq \ker(\psi)}} K_{[\lambda(\psi)]_\mu} \eta^{[\lambda(\psi)]}.$$

Proof. Let $\mathcal{S} = \{\psi \in \text{Irr}(G) \mid L \subseteq \ker(\psi)\}$. For any $m \geq 1$, we have from Proposition 4.4 and Theorem 4.2 that

$$\text{ch}(\text{Ind}_{L_m}^{G_m}(\mathbf{1})) = H_m^{(\mathcal{S})}.$$

Then we have

$$\begin{aligned} \text{Ind}_{L_\mu}^{G_n}(\mathbf{1}) &= \text{Ind}_{L_{\mu_1} \times \dots \times L_{\mu_l}}^{G_n}(\mathbf{1}) \\ &= \text{Ind}_{G_{\mu_1} \times \dots \times G_{\mu_l}}^{G_n} \left(\text{Ind}_{L_{\mu_1}}^{G_{\mu_1}}(\mathbf{1}) \cdots \text{Ind}_{L_{\mu_l}}^{G_{\mu_l}}(\mathbf{1}) \right). \end{aligned}$$

From these two statements, and Theorem 4.2, we have $\text{ch}(\text{Ind}_{L_\mu}^{G_n}(\mathbf{1})) = H_\mu^{(\mathcal{S})}$. The result now follows by applying Lemma 4.3 and Theorem 4.2. □

In the case that $G = L$, Theorem 4.5 boils down to Young’s rule (2.3), and is implied by [23, Proposition 3.12]. In the general setting that $L \trianglelefteq G$ and G/L is abelian, Theorem 4.5 appears to be a new result, stemming exactly from Proposition 4.4, along with Lemma 4.3. Since Lemma 4.3 is implied by the wreath product versions of the Littlewood-Richardson rules appearing in [11, 23], then one should

expect that there is a Littlewood-Richardson generalization of Theorem 4.5. It would be interesting to understand the more general situation when there is no restriction on the subgroup L , and we induce an arbitrary character from L_μ to G_n .

An example of Theorem 4.5 which is of particular interest is the case that G is a cyclic group of order r , which we denote by $C^{(r)}$. In this case $G_n = C^{(r)} \wr S_n$ is the complex reflection group $G(r, 1, n)$. We realize $C^{(r)}$ as the multiplicative group of r th roots of unity, with generator $\zeta = e^{2\sqrt{-1}\pi/r}$, so that $C^{(r)} = \{\zeta^{j-1} \mid 1 \leq j \leq r\}$. We label the irreducible characters of $C^{(r)}$ by $\psi_j(c) = c^{j-1}$, for $1 \leq j \leq r$. Then the irreducible characters of G_n are parameterized by $\mathcal{P}_n[r]$, and we write $\lambda(j)$ for the partition corresponding to $\psi_j \in \text{Irr}(C^{(r)})$. The subgroups of $C^{(r)}$ correspond to positive divisors $d|r$, where $C^{(d)} = \langle \zeta^{r/d} \rangle$ is the subgroup of $C^{(r)}$ of order d . Then we have $C^{(d)} \subseteq \ker(\psi_j)$ if and only if $d|(j-1)$. Proposition 4.4 and Theorem 4.5 then give the following.

Corollary 4.6. *Let $r, n \geq 1$, and let d be a positive divisor of r . Then*

$$\text{Ind}_{C^{(d)} \wr S_n}^{C^{(r)} \wr S_n}(\mathbf{1}) = \sum_{\substack{[\gamma(j)] \in \mathcal{P}_n[r] \\ \text{ht}([\gamma(j)])=1 \\ \gamma(j)=0 \text{ if } d \nmid (j-1)}} \eta^{[\gamma(j)]}.$$

More generally, if $\mu \in \mathcal{P}_n$, then

$$\text{Ind}_{C^{(d)} \wr S_\mu}^{C^{(r)} \wr S_n}(\mathbf{1}) = \sum_{\substack{[\lambda(j)] \in \mathcal{P}_n[r] \\ \lambda(j)=0 \text{ if } d \nmid (j-1)}} K_{[\lambda(j)]\mu} \eta^{[\lambda(j)]}, \text{ and } \text{Ind}_{S_\mu}^{C^{(r)} \wr S_n}(\mathbf{1}) = \sum_{[\lambda(j)] \in \mathcal{P}_n[r]} K_{[\lambda(j)]\mu} \eta^{[\lambda(j)]},$$

where the second case is the result of taking $d = 1$ in the first case.

Several cases of the induced representations in Corollary 4.6 appear as parts of the generalized involution models for the complex reflection groups $G(r, 1, n)$ constructed by Marberg [17]. Namely, $\text{Ind}_{S_n}^{C^{(r)} \wr S_n}(\mathbf{1})$ is the piece of the model corresponding to $k = 0$ for r odd, and $\text{Ind}_{C^{(2)} \wr S_n}^{C^{(r)} \wr S_n}(\mathbf{1})$ corresponds to the $k = \ell = 0$ piece for r even in [17, Theorem 5.6]. This model of the complex reflection groups is obtained independently by Mazorchuk and Stroppel [18], and generalized to projective reflection groups by Caselli and Marberg [3].

5 Multiplicity One for Multipartitions

In this section we prove the main result, which is a generalization of Theorem 3.1 to multipartitions. We begin with a necessary condition for a multipartition Kostka number to be 1.

Lemma 5.1. *Suppose $[\lambda(j)] \in \mathcal{P}_n[r]$ and $\mu \in \mathcal{P}_n$, and let $\tilde{\lambda} \in \mathcal{P}_n$ be as in (2.4). Then $K_{[\lambda(j)]\mu} \geq K_{\tilde{\lambda}\mu}$. In particular, if $K_{[\lambda(j)]\mu} = 1$ then $K_{\tilde{\lambda}\mu} = 1$.*

Proof. From Lemma 2.3, if $K_{\tilde{\lambda}\mu} = 0$, then $K_{[\lambda(j)]\mu} = 0$, so we may suppose that $K_{\tilde{\lambda}\mu} > 0$. Let T be some semistandard Young tableau of shape $\tilde{\lambda}$ and weight μ . As

in Example 2.4, we have seen that from T we can construct a semistandard Young r -multitableau $[T(j)]$ of shape $[\lambda(j)]$ and weight μ . Now suppose T' is a semistandard Young tableau of shape $\tilde{\lambda}$ and weight μ which is distinct from T . Then some column, say the i th column, of T is distinct from column i of T' . Suppose that the entries of column i of T are the entries of column k of $T(j)$ of an r -multitableau $[T(j)]$ of shape $[\lambda(j)]$. Through the process described in Section 2.3, we can construct a semistandard Young r -multitableau $[T'(j)]$ with the entries of column i of T' as the entries of column k of $T'(j)$ (since the order of entries is still preserved, since it was preserved in constructing $[T(j)]$). In particular, $[T(j)]$ and $[T'(j)]$ are distinct. It follows that $K_{[\lambda(j)]\mu} \geq K_{\tilde{\lambda}\mu}$. If $K_{[\lambda(j)]\mu} = 1$, then $K_{\tilde{\lambda}\mu} > 0$ by Lemma 2.3, and so $K_{\tilde{\lambda}\mu} = 1$. \square

The converse of the second statement of Lemma 5.1 is false, which we can see by considering $[\lambda(j)]$ defined by $\lambda(1) = \lambda(2) = (1)$ and $\mu = (1, 1)$. Then $\tilde{\lambda} = (2)$, and $K_{\tilde{\lambda}\mu} = 1$ where $T = \begin{bmatrix} 1 & 2 \end{bmatrix}$ is the unique semistandard Young tableau of shape $\tilde{\lambda}$ and weight μ . However, $[T(j)]$ and $[T'(j)]$ defined by $T(1) = \begin{bmatrix} 1 \end{bmatrix}$, $T(2) = \begin{bmatrix} 2 \end{bmatrix}$, and $T'(1) = \begin{bmatrix} 2 \end{bmatrix}$, $T'(2) = \begin{bmatrix} 1 \end{bmatrix}$, are both semistandard Young 2-multitableaux of shape $[\lambda(j)]$ and weight μ .

The following observation is crucial in the proof of the main result.

Lemma 5.2. *Let $[\lambda(j)] \in \mathcal{P}_n[r]$ and $\mu \in \mathcal{P}_n$ such that $K_{[\lambda(j)]\mu} = 1$. Let $[T(j)]$ be the unique semistandard Young r -multitableau of shape $[\lambda(j)]$ and weight μ . If two distinct $\lambda(j)$ have some column of the same length, then every column of that length in $[T(j)]$ must have identical entries.*

Proof. Suppose that two distinct $\lambda(j)$ have some column of the same length, but two columns of that length do not have identical entries in $[T(j)]$. By Lemma 5.1, $K_{\tilde{\lambda}\mu} = 1$, and let T be the unique semistandard Young tableau of shape $\tilde{\lambda}$ and weight μ . By the process described in Section 2.3, we can construct $[T(j)]$ from T by making each column of T the same as some column of some $T(j)$. So, some two columns of the same length in T do not have the same entries, say columns i and $i + 1$. Suppose $\lambda(j_1)$ and $\lambda(j_2)$ are two distinct partitions of $[\lambda(j)]$ with columns of length the same as columns i and $i + 1$ of $\tilde{\lambda}$. Again from the process from Section 2.3, from the tableau T of shape $\tilde{\lambda}$ and weight μ , we can construct a semistandard Young r -multitableau of shape $[\lambda(j)]$ and weight μ by taking columns of T and making them columns of any individual tableau of shape some $\lambda(j)$, as long as left-to-right order is preserved in individual tableau. This means we can take column i of T and make it a column in a tableau of shape $\lambda(j_1)$ and column $i + 1$ of T a column in a tableau of shape $\lambda(j_2)$, or vice versa. This implies $K_{[\lambda(j)]\mu} > 1$, a contradiction. \square

We may now prove the main theorem. We note that in the bipartition ($r = 2$) case, this is presumably implied by the case of a type B/C Lie algebra in the result of Berenshtein and Zelevinskiĭ [2, Theorem 1.2].

Theorem 5.3. *Let $[\lambda(j)] \in \mathcal{P}_n[r]$ and $\mu \in \mathcal{P}_n$ with $\ell(\mu) = l$. Then $K_{[\lambda(j)]\mu} = 1$ if and only if there exists a choice of indices $0 = i_0 < i_1 < \dots < i_t = l$ such that, for*

each $k = 1, \dots, t$, the r -multipartition $[\lambda(j)^k]$ defined by

$$\lambda(j)^k = (\lambda(j)_{i_{k-1}+1}, \lambda(j)_{i_{k-1}+2}, \dots, \lambda(j)_{i_k}),$$

where $\lambda(j)_i$ is the i th part of $\lambda(j)$ with definition $\lambda(j)_i = 0$ if $i > \ell(\lambda(j))$, and the partition $\mu^k = (\mu_{i_{k-1}+1}, \mu_{i_{k-1}+2}, \dots, \mu_{i_k})$, satisfy the following:

- (1) $\tilde{\lambda}^k \supseteq \mu^k$ (so $|\tilde{\lambda}^k| = |\mu^k|$), and
- (2) for at most one j , either $\lambda(j)_{i_{k-1}+1} = \lambda(j)_{i_{k-1}+2} = \dots = \lambda(j)_{i_k-1} > \lambda(j)_{i_k}$ or $\lambda(j)_{i_{k-1}+1} > \lambda(j)_{i_{k-1}+2} = \lambda(j)_{i_{k-1}+3} = \dots = \lambda(j)_{i_k}$, and for all other j , $\lambda(j)_{i_{k-1}+1} = \lambda(j)_{i_{k-1}+2} = \dots = \lambda(j)_{i_k-1} = \lambda(j)_{i_k}$.

Proof. We first assume that $[\lambda(j)]$ and μ satisfy the listed conditions, and show that $K_{[\lambda(j)]\mu} = 1$. This direction of the proof parallels the same direction of the proof of Theorem 3.1. Since $\tilde{\lambda}^k \supseteq \mu^k$ for $k = 1, \dots, t$, then $\tilde{\lambda} \supseteq \mu$, and so $K_{[\lambda(j)]\mu} > 0$ by Lemma 2.3. Let $[T(j)]$ be some r -multitableau of shape $[\lambda(j)]$ and weight μ . The entries in each column of each $T(j)$ strictly increase, and so the entries in row i of any $T(j)$ must be all at least i . Since row $i_1 + 1$ of each $T(j)$ must have entries at least $i_1 + 1$, then all entries 1 through i_1 must appear in rows 1 through i_1 of $[T(j)]$. Since $|\tilde{\lambda}^1| = |[\lambda(j)^1]| = |\mu^1| = \mu_1 + \dots + \mu_{i_1}$, then also the only entries in rows 1 through i_1 of $[T(j)]$ can be 1 through i_1 . By induction, for each k , the only entries in rows $i_{k-1} + 1$ through i_k of $[T(j)]$ are $i_{k-1} + 1$ through i_k , and these are the only rows in which these entries appear. Thus, we may consider each $[\lambda(j)^k]$ with μ^k independently, and $K_{[\lambda(j)]\mu} = \prod_{k=1}^t K_{[\lambda(j)^k]\mu^k}$. Like in the partition case, the above shows that this factorization holds for any set of indices, as long as $|\tilde{\lambda}^k| = |\mu^k|$ for each k .

It is now enough to show that $K_{[\lambda(j)]\mu} = 1$ whenever $\tilde{\lambda} \supseteq \mu$, $\ell(\mu) = l$, and for at most one j either $\lambda(j)_1 = \lambda(j)_2 = \dots = \lambda(j)_{l-1} > \lambda(j)_l$ or $\lambda(j)_1 > \lambda(j)_2 = \lambda(j)_3 = \dots = \lambda(j)_l$, and for all other j , $\lambda(j)_1 = \lambda(j)_2 = \dots = \lambda(j)_l$. As above, let $[T(j)]$ denote some r -multitableau of shape $[\lambda(j)]$ and weight μ . For all j such that $\lambda(j)_1 = \lambda(j)_2 = \dots = \lambda(j)_l$, then $\ell(\lambda(j)) = l$ (or $\lambda(j) = (0)$), and every column of $T(j)$ must have the entries 1 through l in sequential order (when $\lambda(j) \neq (0)$). If these are the only nonempty $\lambda(j)$, then $[T(j)]$ is uniquely determined and we are done. Otherwise, consider the unique $j = j'$ such that $\lambda(j')_1 > \lambda(j')_2 = \lambda(j')_3 = \dots = \lambda(j')_l$ or $\lambda(j')_1 = \lambda(j')_2 = \dots = \lambda(j')_{l-1} > \lambda(j')_l$. Let s be the total number of columns in all other nonempty $\lambda(j)$, and define the partition ν by $\nu_i = \mu_i - s$. Then $\lambda(j') \supseteq \nu$. Since all other $T(j)$ are uniquely determined, it is enough to show that $K_{\lambda(j')\nu} = 1$. This is implied by Lemma 3.2.

We now assume that $K_{[\lambda(j)]\mu} = 1$. Throughout, let $[T(j)]$ be the unique r -multitableau of shape $[\lambda(j)]$ and weight μ . By Lemma 5.1, we have $K_{\tilde{\lambda}\mu} = 1$. By Theorem 3.1, there exists a choice of indices, say $0 = \iota_0 < \iota_1 < \dots < \iota_u = l$ such that, for each $k = 1, \dots, u$, we have

$$\tilde{\lambda}^k = (\tilde{\lambda}_{\iota_{k-1}+1}, \tilde{\lambda}_{\iota_{k-1}+2}, \dots, \tilde{\lambda}_{\iota_k}) \quad \text{and} \quad \mu^k = (\mu_{\iota_{k-1}+1}, \mu_{\iota_{k-1}+2}, \dots, \mu_{\iota_k}),$$

where $\tilde{\lambda}_i = 0$ if $i > \ell(\tilde{\lambda})$, satisfy $\tilde{\lambda}^k \supseteq \mu^k$ and either

$$\tilde{\lambda}_{\iota_{k-1}+1} = \tilde{\lambda}_{\iota_{k-1}+2} = \dots = \tilde{\lambda}_{\iota_k} \quad \text{or} \quad \tilde{\lambda}_{\iota_{k-1}+1} > \tilde{\lambda}_{\iota_{k-1}+2} = \tilde{\lambda}_{\iota_{k-1}+3} = \dots = \tilde{\lambda}_{\iota_k}.$$

For each $k = 1, \dots, u$, consider the r -multipartition $[\lambda(j)^k]$ given by

$$\lambda(j)^k = (\lambda(j)_{\iota_{k-1}+1}, \lambda(j)_{\iota_{k-1}+2}, \dots, \lambda(j)_{\iota_k}),$$

where $\lambda(j)_\iota = 0$ if $\iota > \ell(\lambda(j))$. Then we have $\tilde{\lambda}^k = \tilde{\lambda}^k$, so that $\tilde{\lambda}^k \supseteq \mu^k$.

Note that whenever $\tilde{\lambda}_i = \tilde{\lambda}_{i+1}$ then $\lambda(j)_i = \lambda(j)_{i+1}$ for each j , since $\lambda(j)_i \geq \lambda(j)_{i+1}$ for each j , and if $\lambda(j)_i > \lambda(j)_{i+1}$ for some j , then $\tilde{\lambda}_i = \sum_j \lambda(j)_i > \sum_j \lambda(j)_{i+1} = \tilde{\lambda}_{i+1}$. Thus, for each $k = 1, \dots, u$, we have

$$\lambda(j)_{\iota_{k-1}+1} = \lambda(j)_{\iota_{k-1}+2} = \dots = \lambda(j)_{\iota_k} \quad \text{or} \quad \lambda(j)_{\iota_{k-1}+2} = \lambda(j)_{\iota_{k-1}+3} = \dots = \lambda(j)_{\iota_k},$$

where either the first holds for every j or the second holds for every j . If for each k , there is at most one j such that either $\lambda(j)_{\iota_{k-1}+1} = \lambda(j)_{\iota_{k-1}+2} = \dots = \lambda(j)_{\iota_k} > \lambda(j)_{\iota_k}$ or $\lambda(j)_{\iota_{k-1}+1} > \lambda(j)_{\iota_{k-1}+2} = \lambda(j)_{\iota_{k-1}+3} = \dots = \lambda(j)_{\iota_k}$, then we can take $t = u$, and $i_k = \iota_k$ for $k = 1, \dots, t$, and the desired conditions are satisfied. Otherwise, we consider the following possibilities.

Consider any k such that there are at least two j such that $\lambda(j)_{\iota_{k-1}+1} > \lambda(j)_{\iota_{k-1}+2}$, or at least two j such that $\lambda(j)_{\iota_{k-1}} > \lambda(j)_{\iota_k}$. We take as a running example $[\lambda(j)]$ with $\lambda(1) = (3^5, 2^2, 1^2)$, $\lambda(2) = (3^5, 2, 1^3)$, and $\lambda(3) = (5, 3^5, 2, 1^2)$, and $\mu = (10^2, 9^3, 7, 5, 3^2)$. In this example, we have the unique r -multitableau $[T(j)]$ of shape $[\lambda(j)]$ and weight μ is given by

$$T(1) = \begin{array}{|c|c|c|} \hline 1 & 1 & 1 \\ \hline 2 & 2 & 2 \\ \hline 3 & 3 & 3 \\ \hline 4 & 4 & 4 \\ \hline 5 & 5 & 5 \\ \hline 6 & 6 & \\ \hline 7 & 7 & \\ \hline 8 & & \\ \hline 9 & & \\ \hline \end{array}, \quad T(2) = \begin{array}{|c|c|c|} \hline 1 & 1 & 1 \\ \hline 2 & 2 & 2 \\ \hline 3 & 3 & 3 \\ \hline 4 & 4 & 4 \\ \hline 5 & 5 & 5 \\ \hline 6 & 6 & \\ \hline 7 & & \\ \hline 8 & & \\ \hline 9 & & \\ \hline \end{array}, \quad T(3) = \begin{array}{|c|c|c|c|c|} \hline 1 & 1 & 1 & 1 & 2 \\ \hline 2 & 2 & 2 & & \\ \hline 3 & 3 & 3 & & \\ \hline 4 & 4 & 4 & & \\ \hline 5 & 5 & 5 & & \\ \hline 6 & 6 & 6 & & \\ \hline 7 & 7 & 7 & & \\ \hline 8 & & & & \\ \hline 9 & & & & \\ \hline \end{array}.$$

Then $\tilde{\lambda} = (11, 9^4, 7, 5, 3^2)$, and $K_{\tilde{\lambda}\mu} = 1$. We may then take as indices satisfying the conditions of Theorem 3.1 to be $\iota_1 = 3$, $\iota_2 = 6$, and $\iota_3 = 9 = \ell(\mu)$, which gives $\tilde{\lambda}^1 = (11, 9^2)$, $\mu^1 = (10^2, 9)$, $\tilde{\lambda}^2 = (9^2, 7) = \mu^2$, and $\tilde{\lambda}^3 = (5, 3^2) = \mu^3$. Suppose first that $\lambda(j)_{\iota_{k-1}+1} > \lambda(j)_{\iota_{k-1}+2}$ for at least two j . In our example, this holds for $k = 3$, where $\lambda(j)_7 > \lambda(j)_8$ for $j = 1, 3$. Then for each such j , there is at least one column of length $\iota_{k-1} + 1$ in $\lambda(j)$ (in the example, $\lambda(1)$ and $\lambda(3)$ each have a column of length 7). By Lemma 5.2, every column of length $\iota_{k-1} + 1$ must be filled identically in each $T(j)$. This means that the right-most $\lambda(j)_{\iota_{k-1}+1} - \lambda(j)_{\iota_{k-1}+2}$ entries of row $\iota_{k-1} + 1$ of $T(j)$ must be identical. Consider those j for which $\lambda(j)_{\iota_{k-1}+2} > 0$. Then $\lambda(j)^k$ has no parts equal to 0 for these j , and so $\ell(\lambda(j)^k) = \ell(\mu^k) = \iota_k - \iota_{k-1}$ for these j . It follows that the left-most $\lambda(j)_{\iota_{k-1}+2}$ columns of $\lambda(j)^k$, in $T(j)$, must contain the

entries $\iota_{k-1} + 1$ through ι_k in sequential order, as there is no other choice. Of these, for the j satisfying $\lambda(j)_{\iota_{k-1}+1} > \lambda(j)_{\iota_{k-1}+2}$, the entries in the rest of row $\iota_{k-1} + 1$ in $T(j)$ must also be $\iota_{k-1} + 1$, since $\mu_{\iota_{k-1}+1} \geq \mu_{\iota_{k-1}+2}$. For those j where $\lambda(j)_{\iota_{k-1}+2} = 0$, since all entries in $T(j)$ of row $\iota_{k-1} + 1$ must be the same, then every entry must also be $\iota_{k-1} + 1$. We have now shown that in this situation, every entry in row $\iota_{k-1} + 1$ must be $\iota_{k-1} + 1$ in every $T(j)$, and no other $\iota_{k-1} + 1$ entries appear. Note that this is indeed the case in the example, where every entry in row 7 of each $T(j)$ is 7. In this case, we define $[\lambda(j)^{k-}], [\lambda(j)^{k+}], \mu^{k-}$, and μ^{k+} by

$$\lambda(j)^{k-} = (\lambda(j)_{\iota_{k-1}+1}), \quad \lambda(j)^{k+} = (\lambda(j)_{\iota_{k-1}+2}, \dots, \lambda(j)_{\iota_k}),$$

$$\mu^{k-} = (\mu_{\iota_{k-1}+1}), \quad \text{and} \quad \mu^{k+} = (\mu_{\iota_{k-1}+2}, \dots, \mu_{\iota_k}),$$

and we define $\iota_{k-} = \iota_{k-1} + 1$ and $\iota_{k+} = \iota_k$. In the example, this means we take $\iota_{3-} = 7, \iota_{3+} = 9$, with

$$\lambda(1)^{3-} = (2) = \lambda(3)^{3-}, \quad \lambda(2)^{3-} = (1), \quad \lambda(1)^{3+} = \lambda(2)^{3+} = \lambda(3)^{3+} = (1^2),$$

$$\mu^{3-} = (5), \quad \text{and} \quad \mu^{3+} = (3^2).$$

Note that for every j , the parts of $\lambda(j)^{k+}$ satisfy $\lambda(j)_{\iota_{k-1}+2} = \dots = \lambda(j)_{\iota_k}$, and we also have $\widetilde{\lambda^{k-}} \supseteq \mu^{k-}$ and $\widetilde{\lambda^{k+}} \supseteq \mu^{k+}$.

Next suppose that $\lambda(j)_{\iota_{k-1}} > \lambda(j)_{\iota_k}$ for at least two j . In the running example, this occurs when $k = 2$, where $\lambda(j)_5 > \lambda(j)_6$ for $j = 1, 2$. The analysis of this case is very similar to the above. We know that there is a column of length $\iota_k - 1$ in at least two $\lambda(j)$ (both $\lambda(1)$ and $\lambda(2)$ have columns of length 5 in the example), and so by Lemma 5.2, every column of this length must be filled identically in every $T(j)$. For those j such that $\lambda(j)_{\iota_k} > 0$, we have $\ell(\lambda(j)^k) = \ell(\mu^k)$, and so the left-most $\lambda(j)_{\iota_k}$ columns of $\lambda(j)^k$ must have the sequential entries $\iota_{k-1} + 1$ through ι_k in $T(j)$. It then follows that the right-most $\lambda(j)_{\iota_{k-1}} - \lambda(j)_{\iota_k}$ columns in $\lambda(j)^k$ must have the sequential entries $\iota_{k-1} + 1$ through $\iota_k - 1$ in every $T(j)$, in order for μ^k to be a partition. This must also then hold for those j where $\lambda(j)_{\iota_k} = 0$. Thus, every entry in row ι_k of every $T(j)$ must be ι_k , and no other ι_k entries appear in any $T(j)$ (which again may be observed to be the case in the example). In this case, we define $[\lambda(j)^{k-}], [\lambda(j)^{k+}], \mu^{k-}$, and μ^{k+} by

$$\lambda(j)^{k-} = (\lambda(j)_{\iota_{k-1}+1}, \lambda(j)_{\iota_{k-1}+2}, \dots, \lambda(j)_{\iota_{k-1}}), \quad \lambda(j)^{k+} = (\lambda(j)_{\iota_k}),$$

$$\mu^{k-} = (\mu_{\iota_{k-1}+1}, \mu_{\iota_{k-1}+2}, \dots, \mu_{\iota_{k-1}}), \quad \text{and} \quad \mu^{k+} = (\mu_{\iota_k}),$$

and we define $\iota_{k-} = \iota_k - 1$ and $\iota_{k+} = \iota_k$. In the example, $\iota_{2-} = 5, \iota_{2+} = 6$,

$$\lambda(1)^{2-} = \lambda(2)^{2-} = \lambda(3)^{2-} = (3^2), \quad \lambda(1)^{2+} = \lambda(2)^{2+} = (2), \quad \lambda(3)^{2+} = (3),$$

$$\mu^{2-} = (9^2), \quad \text{and} \quad \mu^{2+} = (7).$$

For every j , the parts of $\lambda(j)^{k-}$ satisfy $\lambda(j)_{\iota_{k-1}+1} = \dots = \lambda(j)_{\iota_{k-1}}$, and we again have $\widetilde{\lambda^{k-}} \supseteq \mu^{k-}$ and $\widetilde{\lambda^{k+}} \supseteq \mu^{k+}$.

We define $[\lambda(j)^{k^-}]$, $[\lambda(j)^{k^+}]$, μ^{k^-} , μ^{k^+} , ι_{k^-} , and ι_{k^+} as above for every k with the property that there are either at least two j such that $\lambda(j)_{\iota_{k^-}+1} > \lambda(j)_{\iota_{k^-}+2}$ or at least two j such that $\lambda(j)_{\iota_{k^-}} > \lambda(j)_{\iota_{k^-}}$. For all other k , we define $[\lambda(j)^{k^-}] = [\lambda(j)^{k^+}] = [\lambda(j)^k]$, $\mu^{k^-} = \mu^{k^+} = \mu^k$, and $\iota_{k^+} = \iota_{k^-} = \iota_k$. We then define the indices

$$\{i_k \mid k = 1, \dots, t\} = \{\iota_{k^-}, \iota_{k^+} \mid k = 1, \dots, u\},$$

where t is equal to the sum of u and number of k such that there are at least two j such that $\lambda(j)_{\iota_{k^-}+1} > \lambda(j)_{\iota_{k^-}+2}$ or at least two j such that $\lambda(j)_{\iota_{k^-}} > \lambda(j)_{\iota_{k^-}}$. Continuing with the example above, this gives $i_1 = 3$, $i_2 = 5$, $i_3 = 6$, $i_4 = 7$, and $i_5 = 9$. Then $[\lambda(j)^k]$ and μ^k are given by

$$\begin{aligned} \lambda(1)^1 &= \lambda(2)^1 = (3^3), & \lambda(3)^1 &= (5, 3^2), & \mu^1 &= (10^2, 9) \\ \lambda(1)^2 &= \lambda(2)^2 = \lambda(3)^2 = (3^2), & \mu^2 &= (9^2), \\ \lambda(1)^3 &= \lambda(2)^3 = (2), & \lambda(3)^3 &= (3), & \mu^3 &= (7), \\ \lambda(1)^4 &= \lambda(3)^4 = (2), & \lambda(2)^4 &= (2), & \mu^4 &= (5), \\ \lambda(1)^5 &= \lambda(2)^5 = \lambda(3)^5 = (1^2), & \mu^5 &= (3^2). \end{aligned}$$

By construction, we now have that the redefined $[\lambda(j)^k]$ and μ^k for $k = 1, \dots, t$, given by

$$\lambda(j)^k = (\lambda(j)_{i_{k-1}+1}, \lambda(j)_{i_{k-1}+2}, \dots, \lambda(j)_{i_k}),$$

where $\lambda(j)_i = 0$ if $i > \ell(\lambda(j))$, and


$$\mu^k = (\mu_{i_{k-1}+1}, \mu_{i_{k-1}+2}, \dots, \mu_{i_k}),$$

satisfy the required conditions. □

It follows from Theorem 5.3 that for any $[\lambda(j)] \in \mathcal{P}_n[r]$, we have $K_{[\lambda(j)]\tilde{\lambda}} = 1$. We now give a generalization of Corollary 3.4, by classifying which multipartitions $[\lambda(j)]$ satisfy $K_{[\lambda(j)]\mu} = 1$ for only $\mu = \tilde{\lambda}$.

Corollary 5.4. *Let $[\lambda(j)] \in \mathcal{P}_n[r]$. The only $\mu \in \mathcal{P}_n$ satisfying $K_{[\lambda(j)]\mu} = 1$ is $\mu = \tilde{\lambda}$ if and only if for each i either (1) $\tilde{\lambda}_i - \tilde{\lambda}_{i+1} \leq 1$, or (2) there exist at least two j such that $\lambda(j)_i - \lambda(j)_{i+1} \geq 1$ (where we take $\tilde{\lambda}_i = 0$ if $i > \ell(\tilde{\lambda})$ and $\lambda(j)_i = 0$ if $i > \ell(\lambda(j))$).*

Proof. First suppose that there exists some $i = i'$, which we fix, such that $\tilde{\lambda}_{i'} - \tilde{\lambda}_{i'+1} > 1$ and $\lambda(j)_{i'} - \lambda(j)_{i'+1} \geq 1$ for at most one j . This implies that $\lambda(j)_{i'} - \lambda(j)_{i'+1} \geq 1$ for a unique $j = j'$, and so $\tilde{\lambda}_{i'} - \tilde{\lambda}_{i'+1} = \lambda(j')_{i'} - \lambda(j')_{i'+1} > 1$. We will construct a $\mu \in \mathcal{P}_n$ such that $\mu \neq \tilde{\lambda}$ and $K_{[\lambda(j)]\mu} = 1$. Define μ to have parts $\mu_i = \tilde{\lambda}_i$ if $i \neq i', i'+1$, $\mu_{i'} = \tilde{\lambda}_{i'} - 1$, and $\mu_{i'+1} = \tilde{\lambda}_{i'+1} + 1$. Note that μ is a well-defined partition since $\tilde{\lambda}_{i'} - \tilde{\lambda}_{i'+1} \geq 2$. If $\ell(\tilde{\lambda}) = l$, then $\ell(\mu) = l$ by definition. Choose indices as in Theorem



Note that for $i = 1$ and $i = 5$ we have $\tilde{\lambda}_i - \tilde{\lambda}_{i+1} = 0 \leq 1$, while $\tilde{\lambda}_i - \tilde{\lambda}_{i+1} > 1$ for $i = 2, 3, 4, 6$. But, for $j = 1, 2$ we have $\lambda(j)_2 - \lambda(j)_3 = 1$; for $j = 2, 3$ we have $\lambda(j)_3 - \lambda(j)_4 = 1$; for $j = 1, 3$ we have $\lambda(j)_4 - \lambda(j)_5 = 1$; and for $j = 1, 2, 3$ we have $\lambda(j)_6 - \lambda(j)_7 = 1$. That is, for every i either condition (1) or (2) of Corollary 5.4 holds, and thus $K_{[\lambda(j)]\mu} = 1$ only for $\mu = \tilde{\lambda}$.

6 Remarks on Computation and An Application to Finite General Linear Groups

Narayanan considered the computational complexity of evaluating Kostka numbers $K_{\lambda\mu}$ and Littlewood-Richardson coefficients $c_{\nu,\mu}^\lambda$, and showed [19, Proposition 1] that the question of whether $K_{\lambda\mu} > 0$ can be answered in polynomial time. It follows immediately from Lemma 2.3 that the same is true for whether $K_{[\lambda(j)]\mu} > 0$. Further, Ikenmeyer [10, Theorem 3.2] has shown that the question of whether $c_{\nu,\mu}^\lambda$ takes any particular value can be answered in polynomial time. It is likely that the methods of Ikenmeyer translate to the multipartition case, but at the very least it is not difficult to show from Theorem 5.3 that the question of whether $K_{[\lambda(j)]\mu} = 1$ can be answered in polynomial time. Now let \mathbb{F}_q be a finite field with q elements, and let $G = \text{GL}(n, \mathbb{F}_q)$ be the group of invertible n -by- n matrices over \mathbb{F}_q . In this section we define another Kostka number which appears in the representation theory of G , and we see that the computational complexity question for these numbers has quite a different answer than the cases just mentioned.

The complex irreducible characters of $G = \text{GL}(n, \mathbb{F}_q)$ were first described by Green [9]. We give a parameterization of the characters here which follows Macdonald [16, Chapter IV]. If \mathcal{O} is a collection of finite sets, and n is a non-negative integer, then an \mathcal{O} -multipartition of n is a function $\lambda : \mathcal{O} \rightarrow \mathcal{P}$ such that

$$|\lambda| = \sum_{\phi \in \mathcal{O}} |\phi| |\lambda(\phi)| = n.$$

So, an r -multipartition is an \mathcal{O} -multipartition where we take $\mathcal{O} = \{\{1\}, \dots, \{r\}\}$. We let $\mathcal{P}_n^\mathcal{O}$ denote the set of all \mathcal{O} -multipartitions of n , and $\mathcal{P}^\mathcal{O} = \bigcup_{n \geq 0} \mathcal{P}_n^\mathcal{O}$.

Fix an algebraic closure $\bar{\mathbb{F}}_q$ of \mathbb{F}_q , and let $\bar{\mathbb{F}}_q^\times$ be the multiplicative group of $\bar{\mathbb{F}}_q$. Then the Frobenius map F , defined by $F(a) = a^q$, acts on $\bar{\mathbb{F}}_q^\times$, where the set of fixed points of F^m is exactly $\mathbb{F}_{q^m}^\times$, where $m > 0$ is an integer. Let $\hat{\mathbb{F}}_{q^m}^\times$ denote the group of complex characters of $\mathbb{F}_{q^m}^\times$. When $k|m$, there is the standard norm map from $\mathbb{F}_{q^m}^\times$ down to $\mathbb{F}_{q^k}^\times$, which gives rise to the transposed norm map of character groups from $\hat{\mathbb{F}}_{q^k}^\times$ to $\hat{\mathbb{F}}_{q^m}^\times$. We define \mathcal{X} to be the direct limit of the character groups $\hat{\mathbb{F}}_{q^m}^\times$ with respect to these norm maps, $\mathcal{X} = \varinjlim \hat{\mathbb{F}}_{q^m}^\times$.

The set \mathcal{X} is in some sense a dual of $\overline{\mathbb{F}}_q^\times$, by taking consistent (non-canonical) bijections between each $\mathbb{F}_{q^m}^\times$ and its character group. The Frobenius map F acts on \mathcal{X} through the action on $\overline{\mathbb{F}}_q^\times$, and we define Θ to be the collection of F -orbits of \mathcal{X} . The irreducible complex characters of $G = \text{GL}(n, \mathbb{F}_q)$ are parameterized by the set \mathcal{P}_n^Θ of all Θ -multipartitions of n . Given $\lambda \in \mathcal{P}_n^\Theta$, we let $\chi^\lambda \in \text{Irr}(G)$ denote the complex irreducible character of $\text{GL}(n, \mathbb{F}_q)$ to which it corresponds (not to be confused with characters of the symmetric group, which are not used in this section).

Remark. The set of orbits Θ is in cardinality-preserving bijection with the set of F -orbits of $\overline{\mathbb{F}}_q^\times$, which are in turn in bijection with monic non-constant irreducible polynomials over \mathbb{F}_q , where the degree of the polynomial is the same as the cardinality of the corresponding orbit.

Let U denote the group of unipotent upper triangular matrices in G , so

$$U = \{(u_{ij}) \in G \mid u_{ii} = 1, u_{ij} = 0 \text{ if } i > j\}.$$

Fix a non-trivial linear character $\theta : \mathbb{F}_q^+ \rightarrow \mathbb{C}^\times$ from the additive group of \mathbb{F}_q to the multiplicative group of complex numbers. Given any $\mu \in \mathcal{P}_n$, define I_μ to be the complement in $\{1, 2, \dots, n\}$ of the set of partial sums of μ , so

$$I_\mu = \{1, 2, \dots, n\} \setminus \left\{ \sum_{i=1}^j \mu_i \mid 1 \leq j \leq \ell(\mu) \right\}.$$

Define the linear character κ_μ of U by

$$\kappa_\mu((u_{ij})) = \theta \left(\sum_{i \in I_\mu} u_{i, i+1} \right).$$

The *degenerate Gel'fand-Graev character* Γ_μ of G is now defined as

$$\Gamma_\mu = \text{Ind}_U^G(\kappa_\mu).$$

Gel'fand and Graev [8] first considered Γ_μ for $\mu = (n)$, and proved its decomposition into irreducibles is multiplicity free. Zelevinsky [24, Theorem 12.1] decomposed Γ_μ for arbitrary μ as a linear combination of the irreducible characters of G , which in order to describe we must define another variant of the Kostka number. Given $\lambda \in \mathcal{P}_n^\Theta$, let $\Theta_\lambda = \{\varphi \in \Theta \mid \lambda(\varphi) \neq (0)\}$, so Θ_λ is the support of λ . Given $\mu \in \mathcal{P}_n$, a *semistandard Young Θ -multitableau of shape λ and weight μ* , call it \mathbf{T} , is a sequence of semistandard Young tableaux $[\mathbf{T}(\varphi)]$ indexed by $\varphi \in \Theta_\lambda$, such that each $\mathbf{T}(\varphi)$ has shape $\lambda(\varphi)$, and if $\omega(\varphi)$ is the weight of $\mathbf{T}(\varphi)$, then the weight μ of \mathbf{T} has parts

$$\mu_i = \sum_{\varphi \in \Theta_\lambda} |\varphi| \omega(\phi)_i.$$

Then the Kostka number $K_{\lambda\mu}$ is defined to be the total number of semistandard Young Θ -multitableaux of shape λ and weight μ .

Theorem 6.1 (Zelevinsky). *Given any $\mu \in \mathcal{P}_n$ and any $\chi^\lambda \in \text{Irr}(G)$, we have $\langle \Gamma_\mu, \chi^\lambda \rangle = K_{\lambda\mu}$. That is, the degenerate Gel'fand-Graev character Γ_μ has the decomposition*

$$\Gamma_\mu = \sum_{\lambda \in \mathcal{P}_n^\Theta} K_{\lambda\mu} \chi^\lambda.$$

Zelevinsky noted that if μ has parts $\mu_i = \sum_{\varphi \in \Theta} |\varphi| \lambda(\varphi)_i$, then $K_{\lambda\mu} = 1$, and he applied this to show that every irreducible character of G has Schur index 1 [24, Proposition 12.6].

If $\lambda \in \mathcal{P}_n^\Theta$, let $[\lambda(\varphi)]$ denote the $|\Theta_\lambda|$ -multipartition indexed by $\varphi \in \Theta_\lambda$ (where we fix some order of the elements of Θ_λ). So if $|\Theta_\lambda| = r$, then $[\lambda(\varphi)] \in \mathcal{P}[r]$, and we may consider the partition $\tilde{\lambda}$, where $|\tilde{\lambda}| = |[\lambda(\varphi)]|$. If $w > 0$ is an integer, and $\mu \in \mathcal{P}_n$ is such that w divides every part of μ , then we let μ/w denote the partition with parts μ_i/w . Now if $\lambda \in \mathcal{P}_n^\Theta$ has the property that w divides $|\varphi|$ for every $\varphi \in \Theta_\lambda$, then note that we have $|\tilde{\lambda}| = |[\lambda(\varphi)]| = |\mu/w|$. By restricting our attention to those χ^λ such that all $\varphi \in \Theta_\lambda$ have the same size, we may apply our main results Theorem 5.3 and Corollary 5.4 to Theorem 6.1 to obtain the following.

Corollary 6.2. *Let $\mu \in \mathcal{P}_n$, and suppose $\lambda \in \mathcal{P}_n^\Theta$ is such that there exists an integer $w > 0$ such that $|\varphi| = w$ for every $\varphi \in \Theta_\lambda$, and say $|\Theta_\lambda| = r$. Then the following statements hold.*

1. $\langle \Gamma_\mu, \chi^\lambda \rangle = 1$ if and only if $[\lambda(\varphi)] \in \mathcal{P}_{n/w}[r]$ and $\mu/w \in \mathcal{P}_{n/w}$ satisfy the conditions of Theorem 5.3.
2. If $[\lambda(\varphi)] \in \mathcal{P}_{n/w}[r]$ satisfies the conditions of Corollary 5.4, then $\langle \Gamma_\mu, \chi^\lambda \rangle = 1$ if and only if $\mu/w = \tilde{\lambda}$.

The next natural problem is to understand the Kostka numbers $K_{\lambda\mu}$ in the general case, when the orbits $\varphi \in \Theta_\lambda$ have arbitrary sizes. This appears to be a much more difficult problem, which is perhaps reflected when considering complexity. As we see now, even in a simple case, the computational complexity of checking whether $K_{\lambda\mu} > 0$ is in stark contrast with the results mentioned at the beginning of this section. We conclude with the following.

Proposition 6.3. *Let $\lambda \in \mathcal{P}_n^\Theta$ and $\mu \in \mathcal{P}_n$. The problem of determining whether $K_{\lambda\mu} > 0$, even if $\ell(\mu) = 2$ and $\lambda(\varphi) = (1)$ for all $\varphi \in \Theta_\lambda$, is NP-complete.*

Proof. To show that the general problem is NP, consider as a certificate a semistandard Young Θ -multitableau \mathbf{T} of shape λ with entries 1 through $\ell(\mu)$. Then \mathbf{T} has storage size which is polynomial in $\text{size}(\lambda, \mu)$, where $\text{size}(\lambda, \mu)$ is the number of bits needed to input λ and μ (see [20, Section 1]). If $\omega(\varphi)_i$ is the number of i 's in $\mathbf{T}(\varphi)$, to check that \mathbf{T} has weight μ and thus to verify that $K_{\lambda\mu} > 0$, we must check if $\sum_{\varphi \in \Theta_\lambda} |\varphi| \omega(\varphi)_i = \mu_i$ for every i . Since this can be checked in time which is polynomial in $\text{size}(\lambda, \mu)$, the problem is NP.

As remarked above, the orbits in Θ are in bijection with non-constant monic irreducible polynomials over \mathbb{F}_q , where the cardinality of $\varphi \in \Theta$ is the degree of the

corresponding polynomial. It follows that as we increase q and n , the sizes of the orbits in Θ can be any positive integer, and these sizes might occur with multiplicity as large as we like. Now consider the case that $\lambda(\varphi) = (1)$ for all $\varphi \in \Theta_\lambda$ (in which case $\chi^\lambda \in \text{Irr}(G)$ is a *regular semisimple* character), and $\ell(\mu) = 2, \mu = (\mu_1, \mu_2)$. Then deciding whether $K_{\lambda\mu} > 0$ is the same as determining whether the multiset of positive integers $\{|\varphi| \mid \varphi \in \Theta_\lambda\}$ has some sub-multiset which has sum μ_1 . This is exactly the *subset sum* problem, which is known to be *NP*-complete (see [1, Exercise 2.17], for example). Thus to determine whether $K_{\lambda\mu} > 0$ in general is *NP*-complete. \square

Acknowledgments

The authors thank the anonymous referees for very helpful suggestions to improve this paper. The authors also thank Nick Loehr, Nat Thiem, Sami Assaf, Frans Schalekamp, Andreas Stathopoulos, and Arthur Gregory for helpful conversations. The second-named author was supported in part by a grant from the Simons Foundation (#280496).

References

- [1] S. Arora and B. Barak, “Computational complexity, a modern approach”, Cambridge University Press, Cambridge, 2009.
- [2] A. D. Berenshtein and A. V. Zelevinskii, When is the multiplicity of a weight equal to 1?, *Funct. Anal. Appl.* **24** (1990) no. 4, 259–269.
- [3] F. Caselli and E. Marberg, Isomorphisms, automorphisms, and generalized involution models of projective reflection groups, *Israel J. Math.* **199** (2014) no. 1, 433–483.
- [4] T. Ceccherini-Silberstein, F. Scarabotti and F. Tolli, “Representation theory and harmonic analysis of wreath products of finite groups”, London Math. Soc. Lec. Note Series, 410, Cambridge University Press, Cambridge, 2014.
- [5] W. Fulton, “Young tableaux, with applications to representation theory and geometry”, London Math. Soc. Student Texts, 35, Cambridge University Press, Cambridge, 1997.
- [6] W. Fulton and J. Harris, “Representation theory, a first course”, Graduate Texts in Math., 129, Readings in Mathematics, Springer-Verlag, New York, 1991.
- [7] Z. Gates, B. Goldman and C. R. Vinroot, On the number of partition weights with Kostka multiplicity one, *Electron. J. Combin.* **19** (2012) no. 4, Paper 52, 22 pp.
- [8] I. M. Gel’fand and M. I. Graev, Construction of the irreducible representations of simple algebraic groups over a finite field, *Dokl. Akad. Nauk SSSR* **147** (1962), 529–532.
- [9] J. A. Green, The characters of the finite general linear groups, *Trans. Amer. Math. Soc.* **80** (1955) no. 2, 402–447.

- [10] C. Ikenmeyer, Small Littlewood-Richardson coefficients, [arXiv:1209.1521](https://arxiv.org/abs/1209.1521), 2012.
- [11] F. Ingram, N. Jing and E. Stitzinger, Wreath product symmetric functions, *Int. J. Algebra* **3** (2009) no. 1-4, 1–19.
- [12] G.D. James and A. Kerber, “The representation theory of the symmetric group”, Addison-Wesley, Reading, Mass., 1981.
- [13] R. C. King, C. Tollu and F. Toumazet, Stretched Littlewood-Richardson and Kostka coefficients, In: *Symmetry in Physics*, 99–112, CRM Proc. Lec. Notes, 34, Amer. Math. Soc., Providence, RI, 2004.
- [14] A. Knutson and T. Tao, The honeycomb model of $GL_n(\mathbb{C})$ tensor products I: Proof of the saturation conjecture, *J. Amer. Math. Soc.* **12** (1999) no. 4, 1055–1090.
- [15] A. Knutson, T. Tao and C. Woodward, The honeycomb model of $GL_n(\mathbb{C})$ tensor products II: Puzzles determine facets of the Littlewood-Richardson cone, *J. Amer. Math. Soc.* **17** (2004) no. 1, 19–48.
- [16] I. G. Macdonald, “Symmetric functions and Hall polynomials”, second ed., with contributions by A. Zelevinsky, Oxford Math. Monographs, Oxford Science Publications, The Clarendon Press, Oxford University Press, New York, 1995.
- [17] E. Marberg, Generalized involution models for wreath products, *Israel J. Math.* **192** (2012) no. 1, 157–195.
- [18] V. Mazorchuk and C. Stroppel, $G(\ell, k, d)$ -modules via groupoids, *J. Algebraic Combin.* **43** (2016) no. 1, 11–32.
- [19] H. Narayanan, On the complexity of computing Kostka numbers and Littlewood-Richardson coefficients, *J. Algebraic Combin.* **24** (2006) no. 3, 347–354.
- [20] I. Pak and E. Vallejo, Reductions of Young tableau bijections, *SIAM J. Discrete Math.* **24** (2010) no. 1, 113–145.
- [21] B. Sagan, The ubiquitous Young tableau, In: *Invariant theory and tableaux* (Minneapolis, MN, 1988), 262–298; IMA Vol. Math. Appl., 19, Springer, New York, 1990.
- [22] W. Specht, Eine Verallgemeinerung der symmetrischen Gruppe, *Schriften Math. Seminar (Berlin)* **1** (1932), 1–32.
- [23] I. Stein, The Littlewood-Richardson rule for wreath products with symmetric groups and the quiver of the category $F \wr FI_n$, *Comm. Algebra* **45** (2017), 2105–2126.
- [24] A. V. Zelevinsky, “Representations of finite classical groups, a Hopf algebra approach”, Lec. Notes in Math. 869, Springer-Verlag, Berlin-New York, 1981.
- [25] A. V. Zelevinsky, Littlewood-Richardson semigroups, In: *New perspectives in algebraic combinatorics* (Berkeley, CA, 1996-97), 337–345; Math. Sci. Res. Inst. Publ., 38, Cambridge Univ. Press, Cambridge, 1999.

(Received 15 Feb 2016; revised 10 Feb 2017)