

Pairwise additive 1-rotational BIB designs with $\lambda = 1$

KAZUKI MATSUBARA

*Faculty of Commerce, Chuo Gakuin University
Chiba 270-1196
Japan
additivestructure@gmail.com*

SANPEI KAGEYAMA

*Research Center for Math and Science Education
Tokyo University of Science
Tokyo 162-8601
Japan
pkyfc055@ybb.ne.jp*

Abstract

The existence of pairwise additive balanced incomplete block (BIB) designs and pairwise additive cyclic BIB designs with $\lambda = 1$ has been discussed through direct and recursive constructions in the literature. This paper takes BIB designs with 1-rotational automorphisms and then the existence of pairwise additive 1-rotational BIB designs is investigated for $\lambda = 1$. Finally, it is shown that there exists a 2-pairwise additive 1-rotational BIB design with parameters v, k and $\lambda = 1$ if and only if any $v \geq 4$ and $k = 2$.

1 Introduction

A *balanced incomplete block (BIB) design* is a system (V, \mathcal{B}) , where V is a set of v points and \mathcal{B} ($|\mathcal{B}| = b$) is a family of k -subsets (blocks) of V , such that each point of V appears in r different blocks of \mathcal{B} and any two different points of V appear in exactly λ blocks in \mathcal{B} [21]. This is denoted by $\text{BIBD}(v, b, r, k, \lambda)$ or $\text{B}(v, k, \lambda)$.

For a BIB design (V, \mathcal{B}) , let σ be a permutation on V . For a block $B = \{v_1, \dots, v_k\} \in \mathcal{B}$ and a permutation σ on V , let $B^\sigma = \{v_1^\sigma, \dots, v_k^\sigma\}$. When $\mathcal{B} = \{B^\sigma \mid B \in \mathcal{B}\}$, σ is called an *automorphism* of the design (V, \mathcal{B}) . If there exists an automorphism σ of order $v = |V|$, then the BIB design is said to be *cyclic*. On

the other hand, when there exists an automorphism σ of order $v - 1$ with one fixed point, the BIB design is said to be *1-rotational with respect to the cyclic group of order $v - 1$* [2, 19]. Throughout the paper, the BIB design being 1-rotational with respect to the cyclic group of order $v - 1$ is simply said to be 1-rotational. Note that 1-rotational BIB designs with respect to other algebraic groups are not said to be 1-rotational in this paper.

For a cyclic BIB design (V, \mathcal{B}) , the set V of v points can be identified with $Z_v = \{0, 1, \dots, v - 1\}$. In this case, the design has an automorphism $\sigma : i \mapsto i + 1 \pmod{v}$. The *block orbit* containing $B = \{v_1, v_2, \dots, v_k\} \in \mathcal{B}$ is a set of distinct blocks $B + i = \{v_1 + i, v_2 + i, \dots, v_k + i\} \pmod{v}$ for $i \in Z_v$. A block orbit is said to be *full* or *short* according as $|\{B + i \mid 0 \leq i \leq v - 1\}| = v$ or not.

For a 1-rotational BIB design (V, \mathcal{B}) , the set V of v points can be identified with $Z_{v-1} \cup \{\infty\}$ and the block orbit containing $B = \{v_1, v_2, \dots, v_k\} \in \mathcal{B}$ is a set of distinct blocks $B + i = \{v_1 + i, v_2 + i, \dots, v_k + i\} \pmod{v - 1}$ for $i \in Z_{v-1}$. When $B = \{\infty, v_2, \dots, v_k\}$, $B + i = \{\infty, v_2 + i, \dots, v_k + i\}$. Moreover, if $\lambda = 1$, then the orbit of $B = \{\infty, v_2, \dots, v_k\}$ has cardinality $(v - 1)/(k - 1)$. Similarly, a block orbit is said to be full or short according as $|\{B + i \mid 0 \leq i \leq v - 2\}| = v - 1$ or not.

Choose an arbitrary block from each block orbit and call it an *initial block*. The initial block in a full block orbit and a short block orbit is called a full initial block and a short initial block, respectively. It is clear that a cyclic $B(2t, 2, 1)$ or a 1-rotational $B(2t + 1, 2, 1)$ has a short orbit given by the 2-subset $\{0, t\}$ for any $t \geq 1$. This short orbit is denoted by $\{0, t\}\text{PC}(t)$, where $\text{PC}(t)$ means a short cycle of order t , i.e., only $0, 1, \dots, t - 1$ are to be added to the initial block.

Let $s = v/k$, where s need not be an integer, unlike other design parameters. A set of ℓ $\text{BIBD}(v, b, r, k, \lambda)$'s, namely, $(V, \mathcal{B}_1), (V, \mathcal{B}_2), \dots, (V, \mathcal{B}_\ell)$, is called an *ℓ -pairwise additive BIB design*, denoted by $\ell\text{-PAB}(v, k, \lambda)$, if it is possible to pair the designs $(V, \mathcal{B}_1), (V, \mathcal{B}_2), \dots, (V, \mathcal{B}_\ell)$, in such a way that every pair $(V, \mathcal{B}_{i_1}), (V, \mathcal{B}_{i_2})$, where $1 \leq i_1, i_2 \leq \ell$, $i_1 \neq i_2$, gives rise to a new design $(V, \mathcal{B}_{i_1 i_2}^*)$ with parameters $v^* = v = sk$, $b^* = b$, $r^* = 2r$, $k^* = 2k$, $\lambda^* = 2r(2k - 1)/(sk - 1)$. The family $\mathcal{B}_{i_1 i_2}^*$ is defined by $\mathcal{B}_{i_1 i_2}^* = \{B_{i_1 j} \cup B_{i_2 j} \mid 1 \leq j \leq b\}$ with B_{ij} being the j th block of an i th block family \mathcal{B}_i . When $\ell = s$, this is called an *additive BIB design* [16, 23], denoted by $\text{AB}(v, k, \lambda)$. An $\ell\text{-PAB}(v, k, \lambda)$ is said to be cyclic or 1-rotational, denoted by $\ell\text{-PACB}(v, k, \lambda)$ or $\ell\text{-PARB}(v, k, \lambda)$, if (i) every design $(V, \mathcal{B}_1), (V, \mathcal{B}_2), \dots, (V, \mathcal{B}_\ell)$ is cyclic or 1-rotational, respectively, and (ii) every design $(V, \mathcal{B}_{i_1 i_2}^*)$ arising from the pair $(V, \mathcal{B}_{i_1}), (V, \mathcal{B}_{i_2})$ is cyclic or 1-rotational and its initial blocks are obtained by joining an initial block in (V, \mathcal{B}_{i_1}) to an initial block in (V, \mathcal{B}_{i_2}) , where two orbits given by $B_{i_1 j}$ and $B_{i_2 j}$ have the same cardinality for each $1 \leq j \leq b$. Note that if we join an initial block of a $B(v, k, \lambda)$ to an initial block of another $B(v, k, \lambda)$, then the resulting block might not be an initial block of a $B(v, 2k, \lambda')$. When $\ell = s$, this is called an *additive cyclic BIB design* or an *additive 1-rotational BIB design*, denoted by $\text{ACB}(v, k, \lambda)$ or $\text{ARB}(v, k, \lambda)$, respectively. For example, it is checked that the

four block families

$$\begin{aligned}\mathcal{B}_1 & : \{0, 1\}, \{4, 2\}, \{3, 6\}, \{5, \infty\} \pmod{7} \\ \mathcal{B}_2 & : \{4, 2\}, \{0, 1\}, \{5, \infty\}, \{3, 6\} \pmod{7} \\ \mathcal{B}_3 & : \{5, \infty\}, \{3, 6\}, \{0, 1\}, \{4, 2\} \pmod{7} \\ \mathcal{B}_4 & : \{3, 6\}, \{5, \infty\}, \{4, 2\}, \{0, 1\} \pmod{7}\end{aligned}$$

yield an ARB(8, 2, 1). Note that we allow repeated blocks in $(V, \mathcal{B}_{i_1 i_2}^*)$.

Some results on existence are reviewed. In a PAB($v, k, 1$), it is shown that there are an AB($2^n, 2, 1$) and an AB($3^n, 3, 1$) for any integer $n \geq 2$ [22, 23], and there are a 2-PAB($v, 2, 1$) for any $v \geq 4$ and a 3-PAB($v, 2, 1$) for any $v \geq 6$ [11, 14]. Furthermore, partial results on asymptotic existence of ℓ -PAB(v, k, λ)'s and the existence of 2-PACB($v, k, 1$)'s are also shown in [12, 13, 15]. However, for an ℓ -PACB($v, k, 1$), its complete existence is not yet known in the literature, even if $\ell = 2$ and $k = 2$, as the following shows.

Theorem 1.1 [12] There exists a 2-PACB($v, 2, 1$) for any odd integer $v \geq 5$ such that $\gcd(v, 9) \neq 3$.

Theorem 1.2 [15] There exists a 2-PACB($2^m t, 2, 1$) for any integer $m \geq 2$ and any odd integer $t (\geq 1)$ such that $\gcd(t, 27) \neq 3, 9$.

We now focus on 1-rotational BIB designs and the complete existence of a 2-PARB($v, k, 1$) will be established in Section 5 as follows. This will be the main result of the present paper.

Theorem 1.3 There exists a 2-PARB($v, k, 1$) if and only if any $v \geq 4$ and $k = 2$.

Note that the existence of ℓ -pairwise additive BIB designs is equivalent to the existence of some kind of decompositions of a λ -fold complete graph λK_v into edge-disjoint subgraphs isomorphic to a complete graph K_k , denoted by a (v, K_k, λ) -design, in terms of graph embeddings (cf. [3, 7]). In fact, Theorem 1.3 is equivalent to say that there are two 1-rotational $(v, K_2, 1)$ -designs simultaneously embedded into a 1-rotational $(v, K_4, 6)$ -design allowed the repeated blocks such that two K_2 's simultaneously embedded into each K_4 are vertex-disjoint. However, as far as the authors know, any existence result on graphs which is equivalent to Theorem 1.3 has not been provided in literature.

On the other hand, [14] gives a construction of an ℓ -PAB(v, k, λ) by use of nested BIB designs defined in [20]. A survey of nested BIB designs is given in [18] and a more general class of nested BIB designs is further discussed in [10, 17] with wide applicability for other designs. Unfortunately, to the best of our knowledge, by utilizing any result on nested BIB designs we cannot show the complete existence of a 2-PARB($v, 2, 1$).

In particular, Z -cyclic whist tournament designs of order $4n$ in [1] coincide with a special class of nested BIB designs having both a 1-rotational automorphism and

the property of resolvability. It is seen that the Z -cyclic whist tournament designs of order $4n$ can give the $2\text{-PARB}(4n, 2, 1)$ with resolvability by use of the construction method in [14]. However, the investigation of existence of a $2\text{-PARB}(v, 2, 1)$ with resolvability may be as difficult as showing the existence of Z -cyclic whist tournament designs. The resolvability of a $2\text{-PARB}(v, k, 1)$ will be discussed in another paper.

In Section 2, fundamental results for $\text{PAB}(v, k, 1)$'s and 1-rotational BIB designs will be reviewed and the nonexistence of a $2\text{-PARB}(v, k, 1)$ for any $k \geq 3$ will be shown. In Section 3, a pairwise additive cyclic relative difference family (PACDF) used in the proof of Theorem 1.3 will be defined and recursive constructions used in [4, 9, 24] will be developed for the PACDF. Section 4 shows some existence of PACDFs and Section 5 is devoted to the proof of Theorem 1.3. As the appendix, individual examples will be presented.

2 Fundamental results

It is known [23] that in a $\text{PAB}(v, k, \lambda)$

$$2\lambda \equiv 0 \pmod{k-1} \quad (2.1)$$

which implies $k = 2$ or 3 when $\lambda = 1$.

A $\text{B}(v, 3, 1)$ is known as a Steiner triple system (STS). The existence of 1-rotational STSs with respect to an arbitrary group is studied in [2]. Moreover, a characterization of 1-rotational STSs with respect to the cyclic group of order $v - 1$ is known as follows.

Lemma 2.1 [19] Any 1-rotational $\text{B}(v, 3, 1)$ (V, \mathcal{B}) with a point set $V = Z_{v-1} \cup \{\infty\}$ contains the short orbit of the block $\{0, (v-1)/2, \infty\}$ and full orbits in \mathcal{B} .

Now the nonexistence of an $\ell\text{-PARB}(v, k, 1)$ can be shown.

Theorem 2.2 There exists no $\ell\text{-PARB}(v, k, 1)$ for any integers $\ell \geq 2$, $v \geq \ell k$ and $k \geq 3$.

Proof. When $k \geq 4$, (2.1) shows the nonexistence of the design. When $k = 3$, on account of Lemma 2.1, let $\{a, a + (v-1)/2, \infty\}$ and $\{a', a' + (v-1)/2, \infty\}$, $a, a' \in Z_{v-1}$, can be short initial blocks of \mathcal{B}_1 and \mathcal{B}_2 , respectively. By the definition of a $2\text{-PARB}(v, 3, 1)$, \mathcal{B}_{12}^* must contain a set-union of two short initial blocks. However, both of the two blocks contain the element ∞ in common. Hence there does not exist the required design. \square

Remark 2.3 By taking account of an idea used in the proof of Theorem 2.2, a general result can be shown such that there exists no $\ell\text{-PARB}(v, k, (k-1)/2)$ for any $\ell \geq 2$, any $v \geq \ell k$ and any odd integer $k \geq 3$. Hence, it follows from (2.1) that $\lambda \geq k - 1$ in a $\text{PARB}(v, k, \lambda)$.

From now on, we will discuss the remaining case $k = 2$ for $\lambda = 1$ and any $v \geq 4$ to obtain the main result of this paper.

3 Some combinatorial structures

In this section, cyclic difference matrices (CDMs) and cyclic relative difference families (CDFs) are reviewed and pairwise additive cyclic relative difference families (PACDFs) are newly defined. In [4, 9, 24], CDFs are used to construct designs with cyclic (or 1-rotational) automorphisms, and useful recursive constructions of CDFs are given by use of CDMs. Similarly, some constructions of PACDFs are discussed here.

At first CDMs are reviewed. A *cyclic difference matrix* on Z_v , denoted by $\text{CDM}(k, v)$, is defined as a $k \times v$ array $(a(m, n))$, $a(m, n) \in Z_v$, $1 \leq m \leq k$, $1 \leq n \leq v$, that satisfies

$$Z_v = \{a(i, n) - a(j, n) \pmod{v} \mid 1 \leq n \leq v\}$$

for each $1 \leq i < j \leq k$, that is, the differences of any two distinct rows contain every element of Z_v exactly once (see [8]).

Lemma 3.1 [8] There exists a $\text{CDM}(4, v)$ for any odd integer $v \geq 5$ such that $\gcd(v, 27) \neq 9$.

Let G be a group and N be a subgroup of G . Then a family $\mathcal{F} = \{F_i \mid i \in I\}$ of k -subsets of G is called a *relative difference family*, denoted by (G, N, k, λ) -DF, if the list of differences $(d - d' \mid d, d' \in D_i, d \neq d', i \in I)$ contains each element of $G - N$ exactly λ times and each element of N zero time. When G is the cyclic group Z_v and N is the subgroup of Z_v of order n , the relative difference family is said to be *cyclic*, denoted by (v, n, k, λ) -CDF (cf. [4, 24]).

Some results on the existence of $(vg, g, 4, 1)$ -CDFs are known as follows.

Lemma 3.2 [5, 6] There exists a $(2^{s+4}, 2^s, 4, 1)$ -CDF for any integer $s \geq 2$.

Lemma 3.3 [6] There exist a $(81, 9, 4, 1)$ -CDF and a $(243, 27, 4, 1)$ -CDF.

A set of two families \mathcal{F}_1 and \mathcal{F}_2 is called a *2-pairwise additive* (vg, g, k, λ) -CDF, denoted by 2 - (vg, g, k, λ) -PACDF, if both \mathcal{F}_1 and \mathcal{F}_2 are (vg, g, k, λ) -CDFs and the family of set-unions of the j th k -subsets $B_j^{(1)} \in \mathcal{F}_1$ and $B_j^{(2)} \in \mathcal{F}_2$, $1 \leq j \leq |\mathcal{F}_1| = |\mathcal{F}_2|$, is also a $(vg, g, 2k, \lambda')$ -CDF with $\lambda' = 2\lambda(2k-1)/(k-1)$. Throughout the paper, the above “ 2 - (vg, g, k, λ) -PACDF” is simply denoted by “ (vg, g, k, λ) -PACDF”.

Next, some constructions of $(vg, g, 2, 1)$ -PACDFs are provided.

Lemma 3.4 The existence of a $(vg, g, 4, 1)$ -CDF implies the existence of a $(vg, g, 2, 1)$ -PACDF.

Proof. Let 4-subsets of the $(vg, g, 4, 1)$ -CDF on Z_{vg} be

$$\{a_i, b_i, c_i, d_i\}, \quad 1 \leq i \leq \frac{g(v-1)}{12}.$$

Then it is seen that the following families on subsets of Z_{vg} yield the required $(vg, g, 2, 1)$ -PACDF:

$$\begin{aligned} \mathcal{F}_1 & : \{a_i, b_i\}, \{a_i, c_i\}, \{a_i, d_i\}, \{c_i, b_i\}, \{b_i, d_i\}, \{d_i, c_i\} \\ \mathcal{F}_2 & : \{c_i, d_i\}, \{d_i, b_i\}, \{b_i, c_i\}, \{d_i, a_i\}, \{c_i, a_i\}, \{b_i, a_i\} \end{aligned}$$

for $1 \leq i \leq g(v-1)/12$. □

Note that in the proof of Lemma 3.4 the construction of \mathcal{F}_1 and \mathcal{F}_2 is skillful, since an initial subset of the CDF might not arise from the union of initial subsets belonging to families with other parameters.

Lemma 3.5 Let m be a divisor of g . Then the existence of a $(vg, g, 2, 1)$ -PACDF and a $(g, m, 2, 1)$ -PACDF implies the existence of a $(vg, m, 2, 1)$ -PACDF.

Proof. Let $\mathcal{F}_1, \mathcal{F}_2$ and $\mathcal{F}'_1, \mathcal{F}'_2$ be families of a $(vg, g, 2, 1)$ -PACDF and a $(g, m, 2, 1)$ -PACDF, respectively. Then combined families $\mathcal{F}_h^* = \mathcal{F}_h \cup \{\{vx, vy\} \mid \{x, y\} \in \mathcal{F}'_h\}$ on Z_{vg} , $h = 1, 2$, can yield a $(vg, m, 2, 1)$ -PACDF. □

Lemma 3.6 The existence of a $(vg, g, 2, 1)$ -PACDF and a $\text{CDM}(4, v')$ implies the existence of a $(vv'g, v'g, 2, 1)$ -PACDF.

Proof. Let two families of a $(vg, g, 2, 1)$ -PACDF be

$$\mathcal{F}_h : \{x_{hi}, y_{hi}\}$$

for $1 \leq i \leq g(v-1)/2$ and $h = 1, 2$. Further let A be the $\text{CDM}(4, v')$ with $a(m, n)$ as the (m, n) -entry for $1 \leq m \leq 4$ and $1 \leq n \leq v'$. Then, it can be shown that the following two families yield the required $(vv'g, v'g, 2, 1)$ -PACDF on $Z_{vv'g}$:

$$\mathcal{F}_h^* : \{x_{hi} + a(2h-1, n)vg, y_{hi} + a(2h, n)vg\}$$

for $1 \leq i \leq g(v-1)/2, 1 \leq n \leq v'$ and $h = 1, 2$. In fact, let $\{x_{hj}^*, y_{hj}^*\}$ be the j th subset of \mathcal{F}_h^* for $1 \leq j \leq v'g(v-1)/2$ and $h = 1, 2$. Then, by the property of the $\text{CDM}(4, v')$, it can be checked that the multiset of differences arising from the subsets of \mathcal{F}_h^* , $h = 1, 2$, is composed of (i) $\cup_{j=1}^{v'g(v-1)/2} \{\pm(x_{hj}^* - y_{hj}^*)\} = \{\pm(x_{hi} - y_{hi}) + nvg \mid 1 \leq i \leq g(v-1)/2, 0 \leq n \leq v' - 1\}$ containing every element of $Z_{vv'g} - vZ_{vv'g}$ exactly once for each $h = 1, 2$ and (ii) $\cup_{j=1}^{v'g(v-1)/2} \{\pm(x_{1j}^* - x_{2j}^*), \pm(y_{1j}^* - y_{2j}^*), \pm(x_{1j}^* - y_{2j}^*), \pm(y_{1j}^* - x_{2j}^*)\} = \{\pm(x_{1i} - x_{2i} + nvg), \pm(y_{1i} - y_{2i} + nvg), \pm(x_{1i} - y_{2i} + nvg), \pm(y_{1i} - x_{2i} + nvg) \mid 1 \leq i \leq g(v-1)/2, 0 \leq n \leq v' - 1\}$ containing every element of $Z_{vv'g} - vZ_{vv'g}$ exactly four times. Thus it is seen that both \mathcal{F}_1^* and \mathcal{F}_2^* are $(vv'g, v'g, 2, 1)$ -CDFs, and the family of set-unions $\{x_{1j}^*, y_{1j}^*\} \cup \{x_{2j}^*, y_{2j}^*\}, 1 \leq j \leq v'g(v-1)/2$, yields a $(vv'g, v'g, 4, 6)$ -CDF. The proof is complete. □

Note that full initial blocks of a 2-PACB($v, 2, 1$) with no short initial blocks can be considered as a $(v, 1, 2, 1)$ -PACDF. Hence, it is clear that Lemma 3.6 provides a $(vg, g, 2, 1)$ -PACDF, by use of the 2-PACB($v, 2, 1$) with no short initial blocks and a CDM($4, g$).

On the other hand, it is obvious that there does not exist a CDM($4, 2$). Hence, Lemma 3.6 cannot be utilized for the case of $v' = 2$. However, the following recursive construction can be presented.

Lemma 3.7 The existence of a $(vg, g, 2, 1)$ -PACDF implies the existence of a $(2vg, 2g, 2, 1)$ -PACDF.

Proof. Let two families of a $(vg, g, 2, 1)$ -PACDF be

$$\mathcal{F}_h : \{x_{hi}, y_{hi}\}$$

for $1 \leq i \leq g(v - 1)/2$ and $h = 1, 2$. Then, by choosing arbitrary blocks in each orbit of $\{x_{1i}, y_{1i}\} \cup \{x_{2i}, y_{2i}\}$, without loss of generality it can be assumed that $\{x_{1i}, y_{1i}\} = \{0, i\}$.

Now it can be shown that the following two families yield the required $(2vg, 2g, 2, 1)$ -PACDF on Z_{2vg} :

$$\begin{aligned} \mathcal{F}_1^* & : \{x_{1i}, y_{1i}\}, \{x_{2i}, y_{2i} + \delta_i vg\} \\ \mathcal{F}_2^* & : \{x_{2i}, y_{2i} + \delta_i vg\}, \{x_{1i} + vg, y_{1i} + vg\} \end{aligned}$$

for $1 \leq i \leq g(v - 1)/2$, where $\delta_i = 1$ or 0 according as $|y_{2i} - x_{2i}| < vg/2$ or otherwise. In fact, let $\{x_{hj}^*, y_{hj}^*\}$ be the j th subset of \mathcal{F}_h^* for $1 \leq j \leq g(v - 1)$ and $h = 1, 2$. Then the definition of δ_i implies that $\cup_{j=1}^{g(v-1)} \{\pm(x_{hj}^* - y_{hj}^*)\} = \{\pm(x_{1i} - y_{1i}), \pm(x_{2i} - y_{2i} - \delta_i vg) \mid 1 \leq i \leq g(v - 1)/2\}$ contains every element of $Z_{2vg} - vZ_{2vg}$ exactly once for each $h = 1, 2$. Furthermore, it can be checked that $\cup_{j=1}^{g(v-1)} \{\pm(x_{1j}^* - x_{2j}^*), \pm(y_{1j}^* - y_{2j}^*), \pm(x_{1j}^* - y_{2j}^*), \pm(y_{1j}^* - x_{2j}^*)\} = \{\pm(x_{1i} - x_{2i} + nvg), \pm(y_{1i} - y_{2i} + nvg), \pm(x_{1i} - y_{2i} + nvg), \pm(y_{1i} - x_{2i} + nvg) \mid 1 \leq i \leq g(v - 1)/2, 0 \leq n \leq 1\}$ contains every element of $Z_{2vg} - vZ_{2vg}$ exactly four times. Thus it is seen that both \mathcal{F}_1^* and \mathcal{F}_2^* are $(2vg, 2g, 2, 1)$ -CDFs, and the family of set-unions $\{x_{1j}^*, y_{1j}^*\} \cup \{x_{2j}^*, y_{2j}^*\}$, $1 \leq j \leq g(v - 1)$, yields a $(2vg, 2g, 4, 6)$ -CDF. The proof is complete. \square

The results obtained here will be used in the next section.

4 Existence of $(vg, g, 2, 1)$ -PACDFs

In this section, the discussion on existence of $(vg, g, 2, 1)$ -PACDFs is made by use of direct and recursive methods.

Throughout Sections 4 and 5, let P be any odd integer such that $\gcd(P, 6) = 1$ and $P \geq 5$. Then any prime factor of P is not less than 5.

At first, two classes of $(vg, g, 2, 1)$ -PACDFs are produced by use of direct constructions as the following shows.

Lemma 4.1 Let $P \geq 5$ be an odd integer with $\gcd(P, 6) = 1$. Then there exists a $(2P, 2, 2, 1)$ -PACDF.

Proof. Since $\gcd(2, P) = 1$, the following two families on $Z_2 \times Z_P$ can yield the required $(2P, 2, 2, 1)$ -PACDF on Z_{2P} , by corresponding the element j for $0 \leq j \leq 2P - 1$ to (z, w) , where $j \equiv z \pmod{2}$ and $j \equiv w \pmod{P}$:

$$\begin{aligned} \mathcal{F}_1 & : \{(0, 0), (1, a)\}, \{(0, 0), (0, a)\} \\ \mathcal{F}_2 & : \{(1, 2a), (1, 4a)\}, \{(0, 2a), (1, 4a)\} \end{aligned}$$

for any integer a with $1 \leq a \leq (P - 1)/2$. □

Lemma 4.2 Let $P \geq 5$ be an odd integer with $\gcd(P, 6) = 1$. Then there exists a $(3P, 3, 2, 1)$ -PACDF.

Proof. Since $\gcd(3, P) = 1$, the following two families on $Z_3 \times Z_P$ can yield the required $(3P, 3, 2, 1)$ -PACDF on Z_{3P} , by corresponding the element j for $0 \leq j \leq 3P - 1$ to (z, w) , where $j \equiv z \pmod{3}$ and $j \equiv w \pmod{P}$:

$$\begin{aligned} \mathcal{F}_1 & : \{(0, 0), (1, a)\}, \{(0, a'), (0, -a')\} \\ \mathcal{F}_2 & : \{(0, 2a), (1, 3a)\}, \{(1, 2a'), (1, -2a')\} \end{aligned}$$

for any integers a and a' with $1 \leq a \leq P - 1$ and $1 \leq a' \leq (P - 1)/2$. □

Next, some results on the existence of $(vg, g, 2, 1)$ -PACDFs obtained from $(vg, g, 4, 1)$ -CDFs are shown as follows.

Lemma 4.3 There exists a $(2^{4m+n}, 2^n, 2, 1)$ -PACDF for any $n \in \{2, 3, 4, 5\}$ and any positive integer m .

Proof. Lemma 3.4 with the $(2^{4m+n}, 2^{4m+n-4}, 4, 1)$ -CDF obtained by Lemma 3.2 can provide a $(2^{4m+n}, 2^{4m+n-4}, 2, 1)$ -PACDF for any $m \geq 1$ and any $n \in \{2, 3, 4, 5\}$. Hence, for $m = 1$ the result can be shown. Furthermore, by Lemma 3.5 with a $(2^{4(m+1)+n}, 2^{4m+n}, 2, 1)$ -PACDF, the existence of a $(2^{4m+n}, 2^n, 2, 1)$ -PACDF implies the existence of a $(2^{4(m+1)+n}, 2^n, 2, 1)$ -PACDF for $m \geq 1$. Thus, the proof is complete by mathematical induction on m . □

Lemma 4.4 There exist a $(3^n, 9, 2, 1)$ -PACDF and a $(3^{n'}, 3, 2, 1)$ -PACDF for any even integer $n \geq 4$ and any odd integer $n' \geq 3$.

Proof. By applying Lemma 3.4 with the $(81, 9, 4, 1)$ -CDF and the $(243, 27, 4, 1)$ -CDF given in Lemma 3.3, it is shown that there are a $(81, 9, 2, 1)$ -PACDF and a $(243, 27, 2, 1)$ -PACDF. Furthermore, a $(27, 3, 2, 1)$ -PACDF is given in Example A.9. Hence, for any $n \geq 3$, a $(3^n, 3^{n-2}, 2, 1)$ -PACDF can be obtained by applying Lemma 3.6 with the CDM(4, 27) given by Lemma 3.1. Thus, by applying Lemma 3.5 with a $(3^n, 3^{n-2}, 2, 1)$ -PACDF and a $(3^m, 3^{m-2}, 2, 1)$ -PACDF for $3 \leq m \leq n - 2$ repeatedly,

a $(3^n, 9, 2, 1)$ -PACDF and a $(3^{n'}, 3, 2, 1)$ -PACDF can be obtained for any even integer $n \geq 4$ and any odd integer $n' \geq 3$, respectively. \square

Finally, some results on the existence of $(vg, g, 2, 1)$ -PACDFs are shown by use of recursive constructions as follows.

Lemma 4.5 There exist a $(2 \cdot 3^n, 18, 2, 1)$ -PACDF and a $(2 \cdot 3^{n'}, 6, 2, 1)$ -PACDF for any even integer $n \geq 4$ and any odd integer $n' \geq 3$.

Proof. By applying Lemma 3.7 with the $(3^n, 9, 2, 1)$ -PACDF and $(3^{n'}, 3, 2, 1)$ -PACDF obtained by Lemma 4.4, the proof is complete. \square

Lemma 4.6 There exists a $(2^m 3, 2^{m-1}, 2, 1)$ -PACDF for any integer $m \geq 2$.

Proof. By applying Lemma 3.7 with the $(12, 2, 2, 1)$ -PACDF given in Example A.8 repeatedly, the proof is complete. \square

Lemma 4.7 There exists a $(2^m 3^n, 2^m, 2, 1)$ -PACDF for any integers $m \geq 1$ and $n \geq 2$.

Proof. It follows that a family of initial blocks of the 2-PACB $(3^n, 2, 1)$ for $n \geq 2$ obtained by Theorem 1.1 yields a $(3^n, 1, 2, 1)$ -PACDF. Hence, by applying Lemma 3.7 repeatedly, the proof is complete. \square

Lemma 4.8 Let $P \geq 5$ be an odd integer with $\gcd(P, 6) = 1$. Then there exists a $(2Pq, 2q, 2, 1)$ -PACDF for any odd prime $q \geq 5$.

Proof. By applying Lemma 3.6 with the CDM $(4, q)$ for a prime q and the $(2P, 2, 2, 1)$ -PACDF obtained by Lemmas 3.1 and 4.1, respectively, the proof is complete. \square

Lemma 4.9 Let $P \geq 5$ be an odd integer with $\gcd(P, 6) = 1$. Then there exists a $(2^n P, 2^n, 2, 1)$ -PACDF for any positive integer n .

Proof. It follows that a family of initial blocks of the 2-PACB $(P, 2, 1)$ obtained by Theorem 1.1 yields a $(P, 1, 2, 1)$ -PACDF. Hence, by applying Lemma 3.7 repeatedly, the proof is complete. \square

Lemma 4.10 Let $P \geq 5$ be an odd integer with $\gcd(P, 6) = 1$. Then there exists a $(3^n P, P, 2, 1)$ -PACDF for any integer $n \geq 2$.

Proof. For any $n \geq 2$, it follows that a family of initial blocks of the 2-PACB $(3^n, 2, 1)$ obtained by Theorem 1.1 yields a $(3^n, 1, 2, 1)$ -PACDF. Hence, by applying Lemma 3.6 with the CDM $(4, P)$ obtained by Lemma 3.1, the proof is complete. \square

Lemma 4.11 Let $P \geq 5$ be an odd integer with $\gcd(P, 6) = 1$. Then there exists a $(2^m 3P, 2^m 3, 2, 1)$ -PACDF for any positive integer m .

Proof. By applying Lemma 3.7 with the $(3P, 3, 2, 1)$ -PACDF obtained by Lemma 4.2 repeatedly, the proof is complete. \square

Lemma 4.12 Let $P \geq 5$ be an odd integer with $\gcd(P, 6) = 1$. Then there exists a $(2^m 3^n P, 2^m P, 2, 1)$ -PACDF for any integers $m \geq 1$ and $n \geq 2$.

Proof. By applying Lemma 3.7 with the $(3^n P, P, 2, 1)$ -PACDF obtained by Lemma 4.10 repeatedly, the proof is complete. \square

Each of the above-mentioned results will play an important role to show the existence of a 2-PARB($v, 2, 1$) in the next section.

5 Proof of Theorem 1.3

In this section, Theorem 1.3 as the main result of this paper is established. At first a class of 2-PARB($v, 2, 1$)’s is formed.

Lemma 5.1 There exists a 2-PARB($v, 2, 1$) for any $v \geq 6$ with $\gcd(v - 1, 6) = 1$.

Proof. First note that the condition $\gcd(v - 1, 6) = 1$ implies $\{\pm ta \mid 2 \leq a \leq (v - 2)/2\} = Z_{v-1} \setminus \{0, \pm t\}$ on Z_{v-1} for any $t \in \{1, 2, 3\}$. Then, it can be shown that the following block families on $Z_{v-1} \cup \{\infty\}$ yield the required 2-PARB($v, 2, 1$) having

$$\begin{aligned} \mathcal{B}_1 & : \{0, 1\}, \{0, \infty\}, \{0, a\} \pmod{v - 1} \\ \mathcal{B}_2 & : \{2, \infty\}, \{2, 3\}, \{2a, 3a\} \pmod{v - 1} \end{aligned}$$

for any integer a with $2 \leq a \leq (v - 2)/2$. \square

Note that Lemma 5.1 reveals a generalization of Theorem 2.5 in [11], since any odd prime $v - 1$ satisfies $\gcd(v - 1, 6) = 1$.

Next, a class of 2-PARB($v, 2, 1$)’s can be produced as the following shows.

Lemma 5.2 There exists a 2-PARB($2p + 1, 2, 1$) for any odd prime p .

Proof. When $p = 3, 5, 7$, Examples A.2, A.4 and A.6 yield the required designs. Next let $p \geq 11$. Then it can be shown that the following block families yield a 2-PAB($v, 2, 1$) on $Z_2 \times Z_p \cup \{\infty\}$:

$$\begin{aligned} \mathcal{B}_1 & : \{(0, 2), (1, 1)\}, \{(0, 4), (1, 2)\}, \{(0, 0), (1, 3)\}, \{(0, 0), (1, 4)\}, \\ & \quad \{(0, 0), \infty\}, \{(0, 0), (1, a)\}, \{(0, 0), (0, a')\}, \\ & \quad \{(0, 0), (1, 0)\}PC(p) \pmod{(2, p)} \\ \mathcal{B}_2 & : \{(1, 2), (1, 4)\}, \{(1, 4), (1, 8)\}, \{(1, 6), (1, 12)\}, \{(0, 12), \infty\}, \\ & \quad \{(1, 8), (1, 16)\}, \{(1, 2a), (1, 4a)\}, \{(0, 2a'), (1, 4a')\}, \\ & \quad \{(0, 4), (1, 4)\}PC(p) \pmod{(2, p)} \end{aligned}$$

for any integers a and a' with $5 \leq a \leq (p - 1)/2$ and $1 \leq a' \leq (p - 1)/2$. Since $\gcd(2, p) = 1$ implies $Z_2 \times Z_p \cong Z_{2p}$, the required 2-PARB($2p + 1, 2, 1$) on $Z_{2p} \cup \{\infty\}$ can be constructed, by corresponding the element j for $0 \leq j \leq 2p - 1$ to (z, w) , where $j \equiv z \pmod{2}$ and $j \equiv w \pmod{p}$. \square

Next, some results on the existence of a 2-PARB($v, 2, 1$) are shown by use of the observation on $(vg, g, 2, 1)$ -PACDFs given in Section 4 and the following recursive construction.

Lemma 5.3 The existence of a $(vg, g, 2, 1)$ -PACDF and a 2-PARB($g + 1, 2, 1$) implies the existence of a 2-PARB($vg + 1, 2, 1$).

Proof. Let $\mathcal{F}_1, \mathcal{F}_2$ be two families of a $(vg, g, 2, 1)$ -PACDF. Further let two families of initial blocks of a 2-PARB($g + 1, 2, 1$) be

$$\mathcal{F}'_h : \{x_i^{(h)}, y_i^{(h)}\}$$

for $1 \leq i \leq \lfloor (g + 2)/2 \rfloor$ and $h = 1, 2$. Then $\mathcal{F}_h^* = \mathcal{F}_h \cup v\mathcal{F}'_h$, $h = 1, 2$, can yield a 2-PARB($vg + 1, 2, 1$) with

$$v\mathcal{F}'_h : \{vx_i^{(h)}, vy_i^{(h)}\}$$

on $Z_{vg} \cup \{\infty\}$ for $1 \leq i \leq \lfloor (g + 2)/2 \rfloor$ and $h = 1, 2$. \square

The following example illustrates Lemma 5.3 with $v = 9$ and $g = 3$.

Example 5.4 Let \mathcal{F}_1 and \mathcal{F}_2 be two families of the $(27, 3, 2, 1)$ -PACDF given in Example A.9. Furthermore, two families of initial blocks on $Z_{27} \cup \{\infty\}$ obtained from the 2-PARB($4, 2, 1$) given in Example A.1 can be

$$\begin{aligned} 9\mathcal{F}'_1 & : \{0, \infty\}, \{9, 18\} \\ 9\mathcal{F}'_2 & : \{9, 18\}, \{0, \infty\}. \end{aligned}$$

Then combined families $\mathcal{F}_h^* = \mathcal{F}_h \cup 9\mathcal{F}'_h$, $h = 1, 2$, yield a 2-PARB($28, 2, 1$).

For a 2-PACB($v, 2, 1$) with families $\mathcal{B}_1, \mathcal{B}_2$ of blocks, two initial blocks $\{a, a + t\} \in \mathcal{B}_1$ and $\{b, b + t\} \in \mathcal{B}_2$, $a, b, t \in Z_v, t \neq v/2$, such that a set-union of the two initial blocks is an initial block of \mathcal{B}_{12}^* , are now called *friend initial blocks*.

Lemma 5.5 There exists a 2-PARB($2^n + 1, 2, 1$) for any integer $n \geq 2$.

Proof. When $n = 2, 3, 4$, the respective existence of a 2-PACB($2^n, 2, 1$) with friend initial blocks can be seen in [12], i.e., Example 3.4 with $\{0, 1\}, \{2, 3\}$, Example 3.5 with $\{0, 1\}, \{4, 5\}$ and Example 3.9 with $\{0, 7\}, \{5, 12\}$. When $n = 5$, Lemma 3.2 in [15] gives a 2-PACB($2^5, 2, 1$) with friend initial blocks $\{0, 11\}, \{19, 30\}$.

By replacing the friend initial blocks $\{a, a + t\}$ with $\{a, a + t\}$ and $\{a, \infty\}$, and also $\{b, b + t\}$ with $\{b, \infty\}$ and $\{b, b + t\}$, it is shown that there exists a 2-PARB($2^n + 1, 2, 1$) for $n = 2, 3, 4, 5$.

On the other hand, for any $n' \in \{2, 3, 4, 5\}$ and any integer $s \geq 1$, a $(2^{4s+n'}, 2^{n'}, 2, 1)$ -PACDF can be obtained by Lemma 4.3. Hence, by applying Lemma 5.3 with a 2-PARB($2^{n'} + 1, 2, 1$), the proof is complete. \square

Lemma 5.6 There exists a 2-PARB($3^n + 1, 2, 1$) for any positive integer n .

Proof. When $n = 1, 2$, the existence of the required design is given in Examples A.1 and A.3. On the other hand, Lemma 4.4 shows the existence of a $(3^n, 9, 2, 1)$ -PACDF and a $(3^{n'}, 3, 2, 1)$ -PACDF for any even integer $n \geq 4$ and any odd integer $n' \geq 3$, respectively. Hence, based on these PACDFs, by applying Lemma 5.3 with a 2-PARB($10, 2, 1$) and a 2-PARB($4, 2, 1$), the proof is complete. \square

Lemma 5.7 There exists a 2-PARB($2^m 3^n + 1, 2, 1$) for any positive integers m and n .

Proof. When $(m, n) = (1, 1), (2, 1), (1, 2)$, Examples A.2, A.5 and A.7 show the result, respectively.

Let $m = 1$. Then Lemma 4.5 shows the existence of a $(2 \cdot 3^n, 18, 2, 1)$ -PACDF and a $(2 \cdot 3^{n'}, 6, 2, 1)$ -PACDF for any even integer $n \geq 4$ and any odd integer $n' \geq 3$, respectively. Hence, based on these PACDFs, Lemma 5.3 with a 2-PARB($19, 2, 1$) and a 2-PARB($7, 2, 1$) shows the existence of a 2-PARB($2 \cdot 3^n + 1, 2, 1$) for any integer $n \geq 3$.

Let $m \geq 3$ and $n = 1$. Then the $(2^m \cdot 3, 2^{m-1}, 2, 1)$ -PACDF obtained by Lemma 4.6 and the 2-PARB($2^{m-1} + 1, 2, 1$) as in Lemma 5.5 show the existence of a 2-PARB($2^m \cdot 3 + 1, 2, 1$), by applying Lemma 5.3.

Finally, let $m \geq 2, n \geq 2$. Then a 2-PARB($2^m \cdot 3^n + 1, 2, 1$) can be obtained by applying Lemma 5.3 with the $(2^m \cdot 3^n, 2^m, 2, 1)$ -PACDF and the 2-PARB($2^m + 1, 2, 1$) obtained by Lemmas 4.7 and 5.5, respectively. \square

Lemma 5.8 Let $P \geq 5$ be an odd integer with $\gcd(P, 6) = 1$. Then there exists a 2-PARB($2^n P + 1, 2, 1$) for any positive integer n .

Proof. Let $p \geq 5$ be a prime factor of P and $P/p = Q$. Then $Q \geq 1$.

When $n = 1$, Lemma 5.2 itself shows the result for $Q = 1$. Next, for $Q \geq 5$, a $(2P, 2p, 2, 1)$ -PACDF can be obtained by applying Lemma 4.8. Hence, Lemmas 5.2 and 5.3 show the existence of a 2-PARB($2P + 1, 2, 1$).

When $n \geq 2$, a $(2^n P, 2^n, 2, 1)$ -PACDF can be obtained by Lemma 4.9. Hence, the existence of a 2-PARB($2^n + 1, 2, 1$), on account of Lemma 5.5, implies the existence of a 2-PARB($2^n P + 1, 2, 1$). \square

Lemma 5.9 Let $P \geq 5$ be an odd integer with $\gcd(P, 6) = 1$. Then there exists a 2-PARB($3^n P + 1, 2, 1$) for any positive integer n .

Proof. Let $n = 1$. Then the existing $(3P, 3, 2, 1)$ -PACDF obtained by Lemma 4.2 and the 2-PARB($4, 2, 1$) given in Example A.1 show the existence of the required design by Lemma 5.3.

When $n \geq 2$, Lemma 4.10 can provide a $(3^n P, P, 2, 1)$ -PACDF. On the other hand, a 2-PARB($P + 1, 2, 1$) can be given by Theorem 5.1. Hence, Lemma 5.3 can be used to show the existence of a 2-PARB($3^n P + 1, 2, 1$). \square

Lemma 5.10 Let $P \geq 5$ be an odd integer with $\gcd(P, 6) = 1$. Then there exists a 2-PARB($2^m 3^n P + 1, 2, 1$) for any positive integers m and n .

Proof. Let $m \geq 1$ and $n = 1$. Then a $(2^m \cdot 3P, 2^m \cdot 3, 2, 1)$ -PACDF can be given by Lemma 4.11. Furthermore Lemmas 5.3 and 5.7 show the existence of a 2-PARB($2^m \cdot 3P + 1, 2, 1$).

When $m \geq 1$ and $n \geq 2$, a $(2^m \cdot 3^n P, 2^m P, 2, 1)$ -PACDF can be obtained by Lemma 4.12. Hence, Lemmas 5.3 and 5.8 can be used to show the existence of a 2-PARB($2^m \cdot 3^n P + 1, 2, 1$). \square

Finally, the main result is now established as in Theorem 1.3 by taking Theorem 2.2 and Lemmas 5.1 and 5.5 to 5.10.

Proof of Theorem 1.3. When $\gcd(v - 1, 6) = 1$, Lemma 5.1 shows the existence of a 2-PARB($v, 2, 1$). If $\gcd(v - 1, 6) \neq 1$, then $v - 1 = 2^m 3^n$ or $2^m 3^n P$, where $m \geq 0, n \geq 0, (m, n) \neq (0, 0)$ and $P \geq 5$ is any odd integer such that $\gcd(P, 6) = 1$. Then by using Lemmas 5.5 to 5.10 the existence of a 2-PARB($v, 2, 1$) is shown for any $v \geq 4$. This fact with Theorem 2.2 completes the proof. \square

Remark. Some results on the existence of a 2-PACB($v, 2, 1$) are obtained in [12, 15]. Furthermore, some methods of constructing a 2-PARB($v, 2, 1$) given in this paper can be used to construct 2-PACB($v, 2, 1$)’s. As a result, Theorem 1.2 on the existence of 2-PACB designs would be improved. Even so, we cannot show the existence of a 2-PACB($v, 2, 1$) for *any* v . The existence problem of this cyclic type will be discussed in a forthcoming paper.

Appendix

Some individual examples which can be found by use of a computer are presented. Note that each of these examples cannot be given by the construction methods provided in this paper.

Example A.1 An ARB(4, 2, 1) on $Z_3 \cup \{\infty\}$:

$$\begin{aligned} \mathcal{B}_1 & : \{0, \infty\}, \{1, 2\} \pmod 3 \\ \mathcal{B}_2 & : \{1, 2\}, \{0, \infty\} \pmod 3. \end{aligned}$$

Example A.2 A 3-PARB(7, 2, 1) on $Z_6 \cup \{\infty\}$:

$$\begin{aligned} \mathcal{B}_1 & : \{0, \infty\}, \{0, 1\}, \{0, 2\}, \{0, 3\} \text{PC}(3) \pmod 6 \\ \mathcal{B}_2 & : \{1, 3\}, \{2, \infty\}, \{4, 5\}, \{1, 4\} \text{PC}(3) \pmod 6 \\ \mathcal{B}_3 & : \{4, 5\}, \{3, 5\}, \{3, \infty\}, \{2, 5\} \text{PC}(3) \pmod 6. \end{aligned}$$

Example A.3 A 2-PARB(10, 2, 1) on $Z_9 \cup \{\infty\}$:

$$\begin{aligned}\mathcal{B}_1 &: \{0, \infty\}, \{0, 1\}, \{0, 2\}, \{0, 3\}, \{0, 4\} \pmod 9 \\ \mathcal{B}_2 &: \{4, 7\}, \{2, 4\}, \{3, 4\}, \{4, \infty\}, \{3, 7\} \pmod 9.\end{aligned}$$

Example A.4 A 2-PARB(11, 2, 1) on $Z_{10} \cup \{\infty\}$:

$$\begin{aligned}\mathcal{B}_1 &: \{0, \infty\}, \{0, 1\}, \{0, 2\}, \{0, 3\}, \{0, 4\}, \{0, 5\} \text{PC}(5) \pmod{10} \\ \mathcal{B}_2 &: \{4, 7\}, \{7, 9\}, \{8, 9\}, \{4, 8\}, \{5, \infty\}, \{2, 7\} \text{PC}(5) \pmod{10}.\end{aligned}$$

Example A.5 A 2-PARB(13, 2, 1) on $Z_{12} \cup \{\infty\}$:

$$\begin{aligned}\mathcal{B}_1 &: \{0, \infty\}, \{0, 1\}, \{0, 2\}, \{0, 3\}, \{0, 4\}, \{0, 5\}, \\ &\quad \{0, 6\} \text{PC}(6) \pmod{12} \\ \mathcal{B}_2 &: \{6, 8\}, \{2, 11\}, \{6, \infty\}, \{10, 5\}, \{7, 3\}, \{8, 9\}, \\ &\quad \{1, 7\} \text{PC}(6) \pmod{12}.\end{aligned}$$

Example A.6 A 2-PARB(15, 2, 1) on $Z_{14} \cup \{\infty\}$:

$$\begin{aligned}\mathcal{B}_1 &: \{0, \infty\}, \{0, 1\}, \{0, 2\}, \{0, 3\}, \{0, 4\}, \{0, 5\}, \{0, 6\}, \\ &\quad \{0, 7\} \text{PC}(7) \pmod{14} \\ \mathcal{B}_2 &: \{7, 10\}, \{3, 12\}, \{1, 9\}, \{9, 11\}, \{1, \infty\}, \{9, 10\}, \{4, 8\}, \\ &\quad \{6, 13\} \text{PC}(7) \pmod{14}.\end{aligned}$$

Example A.7 A 2-PARB(19, 2, 1) on $Z_{18} \cup \{\infty\}$:

$$\begin{aligned}\mathcal{B}_1 &: \{0, \infty\}, \{0, 1\}, \{0, 2\}, \{0, 3\}, \{0, 4\}, \{0, 5\}, \{0, 6\}, \\ &\quad \{0, 7\}, \{0, 8\}, \{0, 9\} \text{PC}(9) \pmod{18} \\ \mathcal{B}_2 &: \{7, 10\}, \{4, 12\}, \{7, \infty\}, \{9, 16\}, \{1, 5\}, \{9, 10\}, \{2, 8\}, \\ &\quad \{1, 3\}, \{2, 15\}, \{1, 10\} \text{PC}(9) \pmod{18}.\end{aligned}$$

The following examples of PACDFs are used for recursive constructions in Section 4.

Example A.8 A (12, 2, 2, 1)-PACDF:

$$\begin{aligned}\mathcal{F}_1 &: \{0, 1\}, \{0, 2\}, \{0, 3\}, \{0, 4\}, \{0, 5\} \\ \mathcal{F}_2 &: \{5, 10\}, \{7, 11\}, \{2, 4\}, \{8, 11\}, \{2, 3\}.\end{aligned}$$

Example A.9 A (27, 3, 2, 1)-PACDF:

$$\begin{aligned}\mathcal{F}_1 &: \{0, 1\}, \{0, 2\}, \{0, 3\}, \{0, 4\}, \{0, 5\}, \{0, 6\}, \{0, 7\}, \{0, 8\}, \\ &\quad \{0, 10\}, \{0, 11\}, \{0, 12\}, \{0, 13\} \\ \mathcal{F}_2 &: \{6, 7\}, \{7, 26\}, \{8, 13\}, \{2, 17\}, \{17, 24\}, \{1, 7\}, \{19, 23\}, \{14, 24\}, \\ &\quad \{3, 14\}, \{15, 17\}, \{1, 4\}, \{11, 25\}.\end{aligned}$$

Acknowledgments

The authors would like to thank one of the referees for constructive comments through pointing out an error. They make the paper much more readable.

References

- [1] I. Anderson and N. J. Finizio, Some new Z -cyclic whist tournament designs, *Discrete Math.* **293** (2005), 19–28.
- [2] S. Bonvicini, M. Buratti, G. Rinaldi and T. Traetta, Some progress on the existence of 1-rotational Steiner triple systems, *Des. Codes Cryptogr.* **62** (2012), 63–78.
- [3] D. Bryant and S. El-Zanati, Graph decompositions, In: *The CRC Handbook of Combinatorial Designs (2nd ed.)* (Eds.: C. J. Colbourn and J. H. Dinitz), CRC Press, Boca Raton (2007), 477–486.
- [4] M. Buratti, Recursive constructions for difference matrices and relative difference families, *J. Combin. Des.* **6** (1998), 165–182.
- [5] Y. Chang, Some cyclic BIBDs with block size four, *J. Combin. Des.* **12** (2004), 177–183.
- [6] Y. Chang and Y. Miao, Constructions for optimal optical orthogonal codes, *Discrete Math.* **261** (2003), 127–139.
- [7] R. Diestel, *Graph Theory*, 4th ed., Springer-Verlag, 2010.
- [8] G. Ge, On $(g, 4; 1)$ -difference matrices, *Discrete Math.* **301** (2005), 164–174.
- [9] M. Jimbo, Recursive constructions for cyclic BIB designs and their generalizations, *Discrete Math.* **116** (1993), 79–95.
- [10] S. Kageyama and Y. Miao, Nested designs with block size five and subblock size two, *J. Statist. Plann. Inference* **64** (1997), 125–139.
- [11] K. Matsubara and S. Kageyama, The existence of two pairwise additive $\text{BIBD}(v, 2, 1)$ for any v , *J. Stat. Theory Pract.* **7** (2013), 783–790.
- [12] K. Matsubara and S. Kageyama, Some pairwise additive cyclic BIB designs, *Stat. Appl.* **11** (2013), 55–77.
- [13] K. Matsubara and S. Kageyama, The construction of pairwise additive minimal BIB designs with asymptotic results, *Applied Math.* **5** (2014), 2130–2136.

- [14] K. Matsubara and S. Kageyama, The existence of 3 pairwise additive $B(v, 2, 1)$ for any $v \geq 6$, *J. Combin. Math. Combin. Comput.* **95** (2015), 27–32.
- [15] K. Matsubara, M. Hirao and S. Kageyama, The existence of 2 pairwise additive cyclic BIB designs with an even number of points, *Bull. Inst. Combin. Appl.* **75** (2015), 91–94.
- [16] K. Matsubara, M. Sawa, D. Matsumoto, H. Kiyama and S. Kageyama, An addition structure on incidence matrices of a BIB design, *Ars Combin.* **78** (2006), 113–122.
- [17] M. Mishima, Y. Miao, S. Kageyama and M. Jimbo, Constructions of nested directed BIB designs, *Australas. J. Combin.* **18** (1998), 157–172.
- [18] J. P. Morgan, D. A. Preece and D. H. Rees, Nested balanced incomplete block designs, *Discrete Math.* **231** (2001), 351–389.
- [19] K. T. Phelps and A. Rosa, Steiner triple systems with rotational automorphisms, *Discrete Math.* **33** (1981), 57–66.
- [20] D. A. Preece, Nested balanced incomplete block designs, *Biometrika* **54** (1976), 479–486.
- [21] D. Raghavarao, *Constructions and Combinatorial Problems in Design of Experiments*, Dover, New York, 1988.
- [22] M. Sawa, S. Kageyama and M. Jimbo, Compatibility of BIB designs, *Stat. Appl.* **6** (2008), 73–89.
- [23] M. Sawa, K. Matsubara, D. Matsumoto, H. Kiyama and S. Kageyama, The spectrum of additive BIB designs, *J. Combin. Des.* **15** (2007), 235–254.
- [24] J. Yin, Some combinatorial constructions for optical orthogonal codes, *Discrete Math.* **185** (1998), 201–219.

(Received 24 June 2016; revised 29 Dec 2016, 5 Apr 2017)